Growth and morphological characteristics of low- and high-maintenance
Kentucky bluegrass (Poa pratensis L.) cultivars

ISU
1989
B95
C. 3

by

Michael Gene Burt

A Thesis Submitted to the
Graduate Faculty in Partial Fulfillment of the
Requirements for the Degree of
MASTER OF SCIENCE

Major: Horticulture

Approved:

Signatures have been redacted for privacy

Iowa State University
Ames, Iowa

1989

TABLE OF CONTENTS

	Page
INTRODUCTION	1
MATERIALS AND METHODS	12
RESULTS	21
DISCUSSION	44
REFERENCES	47
ACKNOWLEDGMENTS	52

LIST OF TABLES

	Page
Table 1. Concentrations of elements in nutrient solutions	17
Table 2. Quality ratings from the low fertility, unirrigated cultivar evaluation-1987	22
Table 3. Mean squares from the analysis of the 1987 quality rating data from the low fertility, unirrigated cultivar evaluation	23
Table 4. 1988 quality ratings from the low fertility, unirrigated cultivar evaluation	24
Table 5. Mean squares from the analysis of the 1988 quality rating data from the low fertility, unirrigated cultivar evaluation	25
Table 6. Clipping yield data from the low-fertility, unirrigated cultivar evaluation	27
Table 7. Mean squares from the analysis of clipping yield data from the low fertility, unirrigated cultivar evaluation	28
Table 8. Root depth and clipping yield data from the cultivar growth and morphological characterization study	29
Table 9. Mean squares from the analysis of the root depth and clipping yield data from the cultivar growth and morphological characterization study	30
Table 10. Root and shoot growth data from the cultivar growth and morphological characterization study	31
Table 11. Means squares from the analysis of the root and shoot growth data from the cultivar growth and morphological characterization study	32
Table 12. Shoot morphology data from the cultivar growth and morphological characterization study	34
Table 13. Mean squares from the analysis of the shoot morphology data from the cultivar growth and morphological characterization study	35
Table 14. Stomata density data from field grown cultivars	37

Table 15.	Mean squares from the analysis of the stomata density data	38
Table 16.	Cultivar root and shoot growth characteristics when grown under low and high nitrogen (N) levels	40
Table 17.	Mean squares from the analysis of the study on the effects of nitrogen (N) on cultivar root depth and clipping yields	41
Table 18.	Mean squares from the analysis of the study on the effect of nitrogen (N) level on cultivar root and shoot growth characteristics	42

INTRODUCTION

Plant Responses to Unfavorable Environment

Plants possess physiological and morphological adaptations that allow them to survive periods of unfavorable environmental conditions. They can respond to unfavorable conditions in three ways (47). They can cease growth and die, they can continue growth, or they can form resistant organs or structures that go dormant.

Many grasses persist during hot, dry summers due to their ability to go dormant (52). There are many definitions of perennial grass summer dormancy in the literature. Scarth et al. (40), in 1947, defined dormancy as a condition of limited growth. Laude (30), in 1953, defined summer dormancy as a greatly limited, or the complete cessation of, growth brought on by extended periods of summer heat and dryness. Whalley and Davidson (52), in 1968, defined dormancy as a cessation of growth associated with specific physiological changes and the development of structures which enable plants to survive periods of unfavorable environmental conditions. Arcioni et al. (1), in 1984, defined dormancy as a state in which plant growth is temporarily arrested. Ofir and Kerem (38), in 1982, characterized summer dormancy by the development of regeneration buds and geophilic storage organs, followed by the senescence and death of above-ground parts.

Laude (30) studied the summer dormancy response of 20 perennial grasses in the field at Davis, California. He found that seven species, including Poa scabrella [Poa scabrella (Thurb.) Benth. ex Vasey], went

dormant in summer even when supplied with adequate water. Laude studied Poa scabrella in detail and determined that high temperatures and long daylengths were required to bring on the summer dormancy response regardless of soil moisture level. Moderating temperatures and available water were required to break summer dormancy of Poa scabrella. The remaining 13 perennial grass species continued growth as long as water was available, and these species would resume growth after a period of summer dormancy once water became available.

Kentucky bluegrass (Poa pratensis L.) is a perennial grass species that continues growth in regions where adapted when there is available moisture, regardless of daylength and temperature. Prolonged droughts bring on the summer dormancy response of Kentucky bluegrass. Droughts are usually of a severity to bring on a summer dormancy response only during summer months when high ambient temperatures accompany the low levels of available soil water.

Temperature effects

High ambient temperatures increase the rate of transpiration and respiration (29). This can hasten dehydration and the depletion of plant carbohydrates. High temperatures with a radiative load can result in stomata closure. This decreases the cooling effect of transpiration and increases leaf temperature. Stomatal closure and the subsequent rise in leaf temperature decreases the photosynthetic rate, disturbs nitrogen and lipid metabolism, and can injure cell membranes (29).

The optimum temperature for growth of cool-season grasses is 15-24°C (4). Above this temperature range growth declines rapidly, and very high

temperatures will severely injure or kill cool-season grasses. Baker and Jung (3) found the optimum temperature for shoot growth of Kentucky bluegrass to be 21.6°C. They found that at 34.8°C Kentucky bluegrass produced only one-half the shoot growth as that produced at the optimum temperature for growth. There have been many studies on the heat tolerance of Kentucky bluegrass (27,35,48,51) and the performance of Kentucky bluegrass cultivars exposed to high temperatures (17,49,50). Julander (27) found that the Kentucky bluegrasses were killed when exposed to 16 hours of 48°C. Wehner and Watchske (51) found they could completely kill Kentucky bluegrasses at temperatures above 49°C in a water bath, and they found in greenhouse and growth chamber screening tests that cultivar differences in heat tolerance were small.

Drought effects

Kramer (29) defines drought as an environmental stress of sufficient duration to produce a plant water deficit, that in turn causes disturbances in plant physiological processes. Drought is a meteorological and environmental event (33). Plant water deficits always accompany drought, but plant water deficits can also occur in the presence of apparently adequate soil moisture (29). Environmental conditions that can lead to plant water deficits in the presence of adequate soil moisture include: high evaporative demand that can be caused by wind, low humidity, and high radiative load, cold or frozen soils, high salts in soil solution, poor soil aeration, and root injury. Drought induced plant water deficit effects protein metabolism (32). Plant water deficits can inactivate some plant enzymes such as nitrate

reductase and phosphatase, and it can activate other plant enzymes, such as hydrolytic enzymes and some oxidases (32).

According to Kramer (29), plants can escape drought, or plants can tolerate the effects of drought. Plants that escape drought are annual plants that germinate from seed, grow quickly, and produce viable seed before the effects of drought are felt. Many desert ephemerals possess this drought survival strategy. Annual bluegrass (*Poa annua* L.) is an example of a turf-type grass with this adaptation. However, much variability exists in annual bluegrass. Some phenotypes appear as the bunch type, winter annual, true drought escaping types. Others possess a spreading growth habit and a perennial life cycle (4).

Plants can tolerate drought in two ways according to Kramer (29). These plants can postpone, or they can tolerate the effects of dehydration. Dehydration postponement involves morphological and physiological modifications that reduce transpiration and increase absorption. In the grasses, these include adaptations such as decreased shoot surface area, increased shoot canopy resistance to evapotranspiration, responsive stomata, fewer stomata, smaller stomata aperture, deep root systems, and the formation of dormant, vegetative reproductive structures (29). Levitt's term "drought avoidance" is synonymous to drought postponement (32). He states that the principal adaptation of monocots for reducing water loss is the reduction of shoot surface area by leaf folding or rolling, shedding of leaves, and the compact form of foliage and crown. Levitt states that the principal method to increase absorption is a deep, extensive root system. Levitt states that the root to shoot ratio is also important for drought

avoidance. Plants with a high root to shoot ratio need less water absorbed per unit root area to adequately supply the water needs of the shoot system, and the larger root systems of these plants would explore a larger soil volume.

Dehydration tolerance is the ability of a plant to maintain growth at low tissue water potentials (29). There are differences in plant species in their tolerance to dehydration at the molecular level. Kramer postulates that these plant differences are due chiefly to plant variation in cell membrane structure and enzyme activity. An example of plant dehydration tolerance is through osmotic adjustment. Cell turgor is maintained by osmotic adjustment, and cell growth is possible only as long as cell turgor is maintained (32). Osmotic adjustment occurs when tissue cell solutes are concentrated to reduce the cell internal water potential to a point lower than the soil solution. At that point, soil water can flow into the plant down this water potential gradient. Osmotic adjustment has been demonstrated in the monocot crop plants corn (14), wheat (7), and sorghum (25). And, Nus and Hodges (37) demonstrated osmotic adjustment in Kentucky bluegrass.

Plants can possess more than one type of adaptation to survive the dehydrative effects of drought (29). When drought stressed, differences between plant species and differences among cultivars or ecotypes within a plant species in their survival and performance is probably due to a combination of plant adaptations.

There have been many studies on the tolerance and response of cool-season turfgrasses to drought (2,9,34,43) and the response of Kentucky bluegrass cultivars, in particular, to the effects of drought (12,17,34).

These studies show that the Kentucky bluegrass cultivars differ in their response to drought stress, and in their ability to recover from drought induced dormancy. The cultivars that perform well under unirrigated culture in climates where prolonged summer droughts predisposes Kentucky bluegrass to summer dormancy must be better adapted to tolerate the effects of drought through their inherent drought tolerance mechanisms.

Kentucky bluegrass cultivars have been documented to differ in many shoot morphological and growth characteristics, some of which could have an effect on their ability to reduce transpiration. Nittler (36) found Kentucky bluegrass cultivars to differ in growth habit, leaf width, leaf texture, leaf angle, size of stomata, and number of lateral branches formed. Schery (41) found morphological differences in unselected, naturalized ecotypes of Kentucky bluegrass in the midwestern United States. He found differences in growth habit, leaf width, leaf length, and shoot density.

Variability in rooting characteristics could account for differences in soil water absorptive ability. A rapid rate of root elongation, great degree of root proliferation, and large root mass are all drought tolerance mechanisms because each could aid in dehydration postponement.

Varieties within monocot field crop species have been shown to vary in their rooting characteristics. Differences in the rate of root elongation have been documented for wheat varieties (11,19) and sorghum cultivars (26). Drought-resistant, spring wheat varieties had greater early root proliferation (11).

Studies of root systems have also shown root development variation among cultivars of turfgrass species. Differences in root number and

root mass among divergent origin germplasm of St. Augustinegrass [Stenotaphrum secundatum (Walt.) Kuntze] and zoysiagrass (Zoysia spp.) and in root lengths of divergent origin germplasm of St. Augustinegrass, zoysiagrass, and buffalograss [Buchloe dactyloides (Nutt.) Engel.] have been observed (31). Differences in total root mass produced by three bermudagrass [Cynodon dactylon (L.) Pers.] cultivars have been reported (6). The more drought tolerant bermudagrass cultivars produced greater total root mass, and these drought tolerant bermudagrass cultivars had greater total root mass lower in the soil profile. Distinct differences in root mass at certain depths of the rooting profile have been recorded in field grown red fescue (Festuca rubra L.), perennial ryegrass (Lolium perenne L.), and Kentucky bluegrass (5). Of the 22 Kentucky bluegrass cultivars in the study, Barkenta and Captan had consistently low root mass values, and Merion, Fylking, and Campus had consistently high values at the 5-10 cm depth of the root profiles.

Kentucky Bluegrass

Kentucky bluegrass (Poa pratensis L.) is native to temperate Eurasia (8). It was introduced to this country by early settlers, and its inherent ability to rapidly colonize allowed it to spread through much of the open areas of temperate North America (16). There is a great diversity within the Kentucky bluegrasses. This is due to its natural selection over many years and over a wide area subject to much climatic variation (41,44).

Kentucky bluegrass is used extensively in the cool-humid areas of

the United States (4,45). Its use in warmer, drier portions of the country is limited. Kentucky bluegrass can be successfully grown in warm, dry areas if proper cultural practices are followed. Irrigation is required, and fertilization should be applied so that fall growth is maximized and summer stress is minimized (44). Kentucky bluegrass has a great variety of uses (4,45). It is used as ground cover and forage grass in millions of acres of roadsides, pastures, and open areas. It is used as a general purpose turfgrass on lawns, parks, cemeteries, and airfields. Kentucky bluegrass is also used as a high-maintenance sports turf on golf courses and playing fields for a variety of sporting activities.

Nearly all Kentucky bluegrass cultivars have been developed through plant selection rather than through controlled hybridization (4,18,41,44). This is because the Kentucky bluegrasses are highly apomictic- they produce seed without fusion of the male and female gametes. In the early years of cultivar development, plant selections were made from naturalized stands of Kentucky bluegrass (15). In the 1950s and 1960s, there was a push in the turfgrass seed industry to develop cultivars more resistant to common problem diseases and more tolerant to closer mowing and higher fertility levels (10). For this reason many plant selections were made from stands of Kentucky bluegrass subject to specific turfgrass maintenance practices.

Today, the Kentucky bluegrass cultivars can be broadly classified as common and improved types. The common-type Kentucky bluegrasses were not selected and developed for turf-type characteristics. They were developed mainly for high seed production capabilities and early seed

maturation (discussion on the origin of common Kentucky bluegrass cultivars, 25 Feb. 1989. Dr. Doug Brede, Research Director, Jacklin Seed Co.). Common Kentucky bluegrasses are characterized by narrow leaf blades, upright growth habit, low fertility requirement, and susceptibility to Helmithosporium diseases. Improved Kentucky bluegrasses were selected and developed for better turfgrass growth characteristics and disease resistance. Improved Kentucky bluegrasses are characterized by one or more of the following characteristics: dark green color, wide leaves, prostrate growth habit, tolerance to high fertility, and good resistance to Helmithosporium leaf spot and crown rot (16).

1980 National Kentucky bluegrass test

In 1980, the United States Department of Agriculture and the Maryland State Turfgrass Council sponsored a National Kentucky bluegrass test as part of their National Turfgrass Evaluation Program (46). There were 84 cultivars in the test, and each cultivar was sponsored by turfgrass breeders and seed companies. Seed from one seed lot for each cultivar was distributed to 39 locations throughout the United States and Canada. Forty-five trials were established and data were collected from 1981-1985. The purpose of the National Kentucky bluegrass test was to evaluate the sponsored cultivars on a national and local level under diverse environmental conditions and cultural intensities.

There were two trials established at Iowa State University as part of the 1981-1985 National Kentucky bluegrass test. The low-maintenance trial was established in September 1980. This trial receives a September

application of $0.5 \text{ kg N } 100 \text{ m}^{-2} \text{ yr}^{-1}$ and is nonirrigated. The high-maintenance trial was established in August 1981. This trial receives $2 \text{ kg N } 100 \text{ m}^{-2} \text{ yr}^{-1}$ and supplemental irrigation as needed. Both areas are mowed to 5 cm, and each receives annual grass and broadleaf weed control. Data taken on these areas include monthly quality ratings and Helmithosporium leafspot ratings.

The data from Iowa State University has shown that the Kentucky bluegrass cultivars differed in their relative performance in the two trials (20,21,22,23). The cultivars that consistently performed well in the low-maintenance cultivar evaluation tended to perform poorly in the high-maintenance cultivar evaluation relative to the other cultivars in each trial. The Kentucky bluegrass cultivars that performed poorly in the low-maintenance cultivar evaluation tended to perform well in the high-maintenance cultivar evaluation relative to the other cultivars in each trial. All cultivars performed better in the high-maintenance trial than in the low-maintenance trial.

Ten Kentucky bluegrass cultivars were chosen for use in this research project. They were chosen based solely on their performance from 1981-1986 in the low-maintenance Kentucky bluegrass cultivar evaluation at Iowa State University. Five cultivars were chosen that consistently perform well in the low-maintenance trial, and five cultivars were chosen that consistently perform poorly in the low-maintenance trial.

The purpose of this research project was to determine if certain morphological and growth characteristics are common to Kentucky bluegrass cultivars that consistently perform well, and to cultivars that

consistently perform poorly, in the low-maintenance trial at Iowa State University. This was accomplished by closely observing the cultivars in the field, greenhouse, and laboratory. The cultivars were furthered studied in the greenhouse to determine if their root and shoot growth characteristics changed when grown under low and high nitrogen (N) levels.

MATERIALS AND METHODS

Ten Kentucky bluegrass cultivars were chosen based on their performance from 1983 to 1986 in the low-maintenance cultivar evaluation at Iowa State University (20,21,22,23). Five cultivars chosen that consistently perform well in the low-maintenance trial included K3-162, Kenblue, Vantage, South Dakota Common, and S-21. These will be referred to as the "low-maintenance" Kentucky bluegrass cultivars. Five cultivars chosen that consistently perform poorly in the low-maintenance trial included Bonnieblue, Columbia, Lovegreen, A20, and I-13. These cultivars will be referred to as the "high-maintenance" Kentucky bluegrass cultivars. Field, greenhouse, and microscopic studies were performed on these two groups.

Field Studies

In the first stage of the investigation, detailed data were collected on the 10 cultivars from the low-maintenance Kentucky bluegrass cultivar evaluation that had been established in September, 1980. The soil on this site is a Nicollet (fine-loamy, mixed mesic, Aquic Hapludoll) with a pH of 7.2, 10 mg kg⁻¹ phosphorus, 80 mg kg⁻¹ potassium, and 23 g kg⁻¹ organic matter. This area has received an annual application of 0.5 kg nitrogen 100 m⁻² yr⁻¹ and is unirrigated. Annual grasses and broadleaf weeds were controlled annually with common pre- and postemergence herbicides.

Data were collected on turf quality ratings, based on turfgrass color and percent live tissue cover, on the 15 and 30th of each month $\pm$ 2 days,

from May to October in the 1987 and 1988 growing seasons. The ratings were based on a scale of 9 to 1; 9 best, 6 acceptable, and 1 dead or completely dormant turfgrass.

Clipping yields of the 10 cultivars were taken on 1 September, 12 September, and 15 October 1987 with hand shears modified to catch the grass clippings from 580 cm² of the 2.32 m² plots. The clippings were oven-dried at 80°C for 24 hours and recorded in grams of oven-dried clippings m⁻² of plot area.

Greenhouse Studies

A greenhouse study was initiated to determine the growth and morphological characteristics of the 10 cultivars grown under moderate fertility and adequate moisture in a climate controlled research greenhouse. The growing system was a modification of the system reported by Lehman and Engelke (31).

Four mil, clear polyethylene tubing with 3.2 cm outside diameter was sealed on one end and punctured to provide drainage. Four hundred grams of fritted clay (Balcones Absorb-n-Dry) was mixed with 1.75 grams of Osmocote 14-14-14 fertilizer and poured into each tube. The polyethylene tubes were supported by 60-cm, capped, polyvinyl chloride (PVC) pipe, with 4.3 cm outside diameter. The PVC pipe was supported across the greenhouse bench by a wooden frame and wire fencing. The tubes were vibrated and thoroughly leached with distilled water leaving a 60-cm rooting profile. Mortite (Mortell Company, P.O. Box 71 550 N. Hobbie Ave., Kankakee Ill. 60901) was wrapped around the outside of the polyethylene tube at the top

of the PVC pipe to inhibit light penetration.

Seed of each cultivar was germinated in fritted clay. Individual, uniform, 3-week old seedlings were transplanted to the tubes. The tubes were arranged in a randomized complete block design with four replications across the greenhouse bench. The PVC pipes were angled 30° from vertical to force root growth down the lower side of the tubes which allowed visual monitoring and measurement of root growth.

The tubes were watered daily with distilled water until each seedling was well established. The tubes were then watered weekly to saturation. Because of insufficient natural sunlight intensity and duration, high pressure sodium lamps were used for supplemental lighting during the last four weeks of the experiment. The average light intensity across the greenhouse bench was $675 \mu\text{Mol sec}^{-1} \text{ m}^{-2}$, and the photoperiod was 14 hours. The air temperature was maintained at $25 \pm 2^\circ\text{C}$. Data taken weekly during the course of the experiment included fresh clipping weights and rooting depths. Root depth was measured from the plant crown to the point of maximum root penetration. The plants were mowed at 6.4 cm, and the fresh clipping weights were recorded.

All plant material was harvested 65 days after the seedlings were transplanted into the tubes. At harvest, the final lateral branch number and shoot fresh weights were recorded. The polyethylene rooting columns were divided into three, 20-cm sections, and the fritted clay was carefully and thoroughly washed away from the root tissue. The roots and shoots were oven dried at 80°C for 24 hours, and shoot and root dry weights were recorded.

Shoot morphological data were based on an average of at least five

representative shoots per tube. This data were taken just before or at harvest, and it included leaf width, leaf folding, sheath length, leaf angle, and leaf number per shoot. Leaf width was measured at the widest point of the most recently emerged, fully developed, unmowed leaf blade. Sheath lengths were measured from the crown to the uppermost portion of each sheath. Leaf folding data were taken as an overall rating of all shoots in each tube just before harvest. The ratings were based on a scale of 1 to 5; with 1 being very tightly folded, and 5 being virtually flat leaf blades. Leaf angle from horizontal was measured from the vertical youngest leaf blade to the next youngest leaf blade. The average number of leaves of several representative shoots per tube were also recorded.

The study was conducted three times. The second study was initiated in December 1987, and the third study was initiated in January 1988. High- pressure sodium lamps were used once seedlings were well established in each experiment. A greenhouse cooling system malfunction resulted in poor temperature control during the latter portions of the second study, and during much of the third study. Afternoon temperatures as high as 38°C were recorded. Because of the very warm midday temperatures, the high pressure sodium lamps were used each day only from 6:00 am to 9:00 am, and again from 5:30 pm to 8:30 pm during the months of March and April for each experiment. The second study was harvested at 75 days, and the third study was harvested at 70 days, after the seedlings were transplanted to the polyethylene tubes.

Another greenhouse study was initiated in June, 1988 to determine how both low and high levels of nitrogen (N) affected the growth

characteristics of two low-maintenance and two high-maintenance Kentucky bluegrass cultivars. The low-maintenance cultivars included Vantage and K3-162, and the high-maintenance cultivars included Bonnieblue and I-13.

The same polyethylene tube growth system was used for this study. The rooting media included 900 g of a fine, washed sand (100% passed through a 1.5-mm sieve) per tube. The tubes were vibrated and leached with distilled water leaving a rooting profile of 60 cm.

Plant nutrients for this growth system were provided by nutrient solutions similar to those described by Pellet and Roberts (39). High N (102 ppm) and low N (11 ppm) nutrient solutions were utilized. Each contained high phosphorous (P at 25 ppm) and potassium (K at 63 ppm) levels, and the recommended amounts of the remaining essential nutrients (Table 1).

Uniform, individual, 4-week old seedlings for each cultivar were transplanted to two adjacent tubes per replication in the study. This study was conducted in a split plot design, with cultivars as the whole plot and N levels as the split plot. One tube for each cultivar was saturated daily with the high N, and the other tube for each cultivar was saturated daily with the low N nutrient solution in each of the four replications of the study. The cultivars were assigned randomly in each replication, and the high and low N solutions were also assigned randomly to the paired tubes of each cultivar in each replication. No supplemental lighting was provided, and greenhouse air temperature was maintained at $27 \pm 3^{\circ}\text{C}$.

Table 1. Concentrations of elements in nutrient solutions

Element	p.p.m
High nitrogen	102
Low nitrogen	11
High phosphorus	25
Low phosphorus	4
High potassium	63
Low potassium	10
Calcium	104
Magnesium	19
Sulfur	72-153
Iron	1.2
Boron	0.1
Manganese	0.25
Zinc	0.1
Copper	0.01
Molybdenum	0.01

Rooting depth data were taken weekly during the course of the experiment. Due to the speed at which the roots penetrated the sand media, the rooting rate data were presented in number of days for the roots to penetrate to the bottom of the rooting tubes.

The plant material was harvested 105 days after seedlings were transplanted. At harvest, the final shoot weights and lateral branch numbers were recorded. The polyethylene rooting columns were divided into three, 20-cm sections, and the root tissue was separated from the sand media. The roots and shoots were oven-dried at 80°C for 24 hours, and shoot and root dry weights were recorded.

The study was repeated in July, 1988. Plant material from this study was harvested 110 days after seedlings were transplanted into the tubes.

Laboratory studies

The laboratory data for this investigation included microscopic stomata density counts. Adaxial and abaxial stomatal counts were taken from samples of the 10 Kentucky bluegrass cultivars taken from the high-maintenance Kentucky bluegrass cultivar field evaluation at Iowa State University. The plant material was harvested on 21 June and again on 19 July 1988 from the high-maintenance trial.

Stomatal impressions were made in nitrocellulose using a technique reported by Shearman and Beard (42). The plant material used for stomatal impressions were uniform, fully expanded, unmowed, most recently emerged leaf blades of randomly chosen shoots. The nitrocellulose was painted on the leaf blades and left to thoroughly dry. The nitrocellulose layer was

peeled from the leaf blade and mounted on slides for microscopic observation.

Stomatal density counts were made at 500 magnifications from the midportion of the leaf blades. Five counts were made per leaf blade on both the adaxial and abaxial surfaces. Stomatal density counts were made from four leaves of separate shoots per cultivar. The actual leaf area observed was 0.126 mm^2 , and counts were converted to number of stomata mm^{-2} .

Analysis of data

Data from each of the experiments within the growth and morphology characterization studies, the stomatal density count studies, and the cultivar growth studies under low and high nitrogen (N) levels were analyzed separately, and the data from the experiments within each study were combined for analysis. The field quality rating data for 1987 and 1988 were not combined for analysis, and the clipping yield data from the low fertility, unirrigated trial were also not combined for analysis.

Data from all studies were tested for differences among cultivars. When significant cultivar differences occurred at the 5% level of significance, the cultivar sums of squares was subdivided into single degree of freedom contrasts to test for differences between the combined means of the low-maintenance versus the high-maintenance cultivars. The cultivar sums of squares were further subdivided in the field studies, the growth and morphology characterization study, and the stomatal density count study to assess the variability both among the low-maintenance and

among the high-maintenance cultivars.

Data from the cultivar growth study under low and high N level were also analyzed to determine the effect of the N level, and the N * cultivar interaction on the variability within these data. The N * cultivar interaction was subdivided to determine if the differences between the low-maintenance versus the high-maintenance cultivars changed over N level.

RESULTS

Field Studies

Cultivar quality ratings

During the 1987 growing season, there were no differences between the low- and high-maintenance cultivars during May and June (Tables 2 and 3). From mid-July through the end of October, the low-maintenance cultivars received higher quality ratings. The low-maintenance cultivars were also rated higher for the average over the entire growing season.

Differences occurred among the high-maintenance cultivars on July 15 and July 30. Differences occurred among the low-maintenance cultivars on July 15, July 30, August 15, and for the growing season average. The variability among the low-maintenance cultivars was mostly due to a low rating of Vantage, and the variability among the high-maintenance cultivars was largely due to a low rating of I-13 relative to the other cultivars in the study.

During the 1988 growing season, the high-maintenance cultivars were rated higher on May 15 and June 15 (Tables 4 and 5). The low-maintenance cultivars were rated higher on July 30, August 15, and September 15. When the data were averaged over the entire growing season, there was no difference between the combined means of the low- versus the high-maintenance cultivars.

There were differences among the low-maintenance cultivars on June 15 and July 30, and among the high-maintenance cultivars on July 30 and August 15. Again, the difference among the low-maintenance and among the

Table 2. Quality rating data from the low fertility, unirrigated cultivar field evaluation-1987

Cultivars	Rating date												Season mean
	<u>May</u>		<u>June</u>		<u>July</u>		<u>Aug.</u>		<u>Sept.</u>		<u>Oct.</u>		
	15	30	15	30	15	30	15	30	15	30	15	30	
	Rating†												
	<u>Low-maintenance cultivars</u>												
S-21	4.7	4.7	4.7	4.3	5.3	5.7	5.7	5.7	5.0	5.7	5.0	5.7	5.2
S. D. Common	4.3	4.7	4.3	4.0	5.7	6.0	5.7	5.7	6.0	6.0	6.3	6.0	5.4
K3-162	5.3	5.7	5.0	4.3	6.0	6.0	6.0	6.0	6.3	6.0	5.3	6.0	5.7
Kenblue	3.7	3.6	4.3	4.3	6.0	6.0	6.3	6.3	5.7	6.0	5.3	6.3	5.3
Vantage	4.3	4.3	4.3	4.0	4.3	4.3	4.3	4.7	4.3	5.0	4.7	5.7	4.5
Mean	4.5	4.6	4.5	4.2	5.5	5.6	5.6	5.7	5.5	5.7	5.3	5.9	5.2
	<u>High-maintenance cultivars</u>												
I-13	4.7	5.3	4.7	4.7	2.7	2.3	2.3	2.3	2.7	3.3	3.7	4.7	3.6
Bonnieblue	4.7	5.0	4.7	4.0	4.0	3.7	3.7	3.7	3.3	3.7	4.0	4.0	4.0
A20	3.7	4.0	3.3	3.3	3.7	3.7	3.7	3.3	3.0	3.7	3.7	4.0	3.6
Columbia	4.0	4.0	4.0	4.0	3.7	3.3	3.7	3.0	3.7	4.0	3.7	4.0	3.6
Lovegreen	4.7	5.0	4.7	4.3	4.3	4.3	3.3	2.7	3.3	4.0	3.7	4.0	4.0
Mean	4.3	4.7	4.3	4.1	3.7	3.5	3.3	3.0	3.2	3.7	3.7	4.1	3.8
LSD (0.05)	1.1	1.7	1.1	1.3	0.8	0.8	1.1	1.5	2.1	1.4	1.2	1.4	0.6

[†]Quality rating based on a scale of 9 to 1; 9 = best, 6 = acceptable, 1 - dead or completely dormant turf.

Table 3. Mean squares from the analysis of the 1987 quality rating data from the low fertility, unirrigated cultivar field evaluation

		Rating date												
		May		June		July		Aug.		Sept.		Oct.		Season
Source	df	15	30	15	30	15	30	15	30	15	30	15	30	Average
Replication	2	3.10**	2.23	0.30	0.23	1.23*	0.93	2.43**	3.03*	3.43	2.43*	7.23**	7.64**	2.06**
Cultivar	9	0.80	1.22	0.65	0.38	3.86**	5.20**	5.50**	6.81**	5.33**	3.69**	2.68**	2.92**	1.98**
Contrast†	1	NA†	NA	NA	NA	24.30**	34.13**	38.53**	53.33**	38.53**	30.00**	19.20**	24.30**	15.05**
Among high‡	4	NA	NA	NA	NA	1.00*	1.21*	1.01	0.84	0.43	0.26	0.07	0.30	0.15
Among low§	4	NA	NA	NA	NA	1.44**	1.58**	1.74*	1.11	1.94	0.57	1.09	0.20	0.55*
Pooled error	18	0.43	0.97	0.37	0.53	0.23	0.27	0.40	0.74	1.43	0.66	0.49	0.63	0.14
Corrected total	29													

*, **Significant at the 5% or 1% level, respectively.

†Tests for differences between the combined means of the low- versus the high-maintenance cultivars.

‡, §Tests for differences among the high-maintenance and among the low-maintenance cultivars, respectively.

‡NA indicates contrasts and tests among high and among low not performed where cultivar differences were nonsignificant.

Table 4. Quality rating data from the low fertility, unirrigated cultivar field evaluation-1988

	Rating date-1988												
	<u>May</u>		<u>June</u>		<u>July</u>		<u>Aug.</u>		<u>Sept.</u>		<u>Oct.</u>		Season
Cultivars	15	30	15	30	15	30	15	30	15	30	15	30	mean
	Rating†												
	<u>Low-maintenance cultivars</u>												
S-21	2.3	3.3	2.0	1.0	1.0	3.7	3.0	2.3	3.0	3.0	3.0	3.3	2.6
S. D. Common	2.3	3.3	2.0	1.0	1.0	3.7	3.3	3.0	2.7	3.0	3.0	3.3	2.6
K3-162	2.3	4.7	3.0	1.0	1.0	3.7	3.0	2.3	4.0	4.0	4.0	4.3	3.1
Kenblue	2.3	4.7	2.7	1.0	1.0	4.0	3.3	3.3	3.3	3.3	3.3	3.7	3.0
Vantage	3.7	5.7	4.0	1.0	1.0	5.0	4.0	3.3	4.3	4.7	4.7	5.7	3.9
Mean	2.6	4.3	2.7	1.0	1.0	4.0	3.3	2.9	3.5	3.6	3.6	4.1	3.1
	<u>High-maintenance cultivars</u>												
I-13	4.7	3.3	4.3	1.0	1.0	2.3	2.0	1.7	2.0	1.7	1.7	2.3	2.3
Bonnieblue	3.3	4.0	3.7	1.0	1.0	3.3	3.0	2.3	3.3	3.7	4.0	4.3	3.1
A20	3.3	2.7	3.3	1.0	1.0	4.0	3.3	3.0	3.3	3.3	3.3	3.7	2.9
Columbia	3.3	4.0	3.7	1.0	1.0	2.7	2.7	2.3	2.3	2.7	2.7	3.0	2.6
Lovegreen	3.7	3.3	4.0	1.0	1.0	3.7	3.0	2.7	3.3	3.7	3.7	4.0	3.1
Mean	3.6	3.5	3.8	1.0	1.0	3.2	2.8	2.4	2.9	3.0	3.1	3.5	2.8
LSD (0.05)	1.4	2.1	1.0	0.0	0.0	1.0	0.8	1.0	1.2	2.0	2.2	2.0	0.6

[†]Quality ratings based on a scale of 9 to 1; 9 = best, 6 = acceptable, and 1 = dead or completely dormant turf.

Table 5. Mean squares from the analysis of the 1988 quality rating data from the low fertility, unirrigated cultivar field evaluation

		Rating date												
Source	df	<u>May</u>		<u>June</u>		<u>July</u>		<u>Aug.</u>		<u>Sept.</u>		<u>Oct.</u>		Season average
		15	30	15	30 [¶]	15 [#]	30	15	30	15	30	15	30	
Replication	2	0.53	0.43	0.43	-	-	1.20	1.23*	0.43	2.03*	3.10	3.74	4.23	0.55*
Cultivar	9	1.87*	2.37	2.06**	-	-	1.61**	0.80*	0.85	1.50*	1.96	2.07	2.45	0.57**
Contrast [†]	1	8.53**	NA ^{††}	8.53**	-	-	4.80**	2.13**	NA	2.70*	NA	NA	NA	0.43
Among high [‡]	4	1.01	NA	0.43	-	-	1.44*	0.76*	NA	1.26	NA	NA	NA	0.36
Among low [§]	4	0.97	NA	1.13*	-	-	1.21*	0.50	NA	1.42	NA	NA	NA	0.86**
Pooled error	18	0.64	1.49	0.36	0	0	0.35	0.23	0.36	0.48	1.36	1.59	1.38	0.14
Corrected total	29													

^{*}, ^{**}Significant at the 5% or 1% level, respectively.

[†]Tests for differences between the combined means of the low- versus the high-maintenance cultivars.

[‡], [§]Tests for differences among the high-maintenance and among the low-maintenance cultivars, respectively.

[¶], [#]All cultivars completely dormant on 30 June and 15 July 1988. Ratings for all cultivars=1.

^{††}NA indicates contrasts and tests among high and among low not performed where cultivar differences were nonsignificant.

high-maintenance cultivars was largely due to outlying values for Vantage and I-13. In this case, Vantage was rated significantly higher and, again, I-13 rated lower than the other cultivars in the study.

Cultivar clipping yields

The low-maintenance cultivars produced greater clipping yields on all dates (Tables 6 and 7). No significant variation was observed among low-maintenance or among high-maintenance cultivars in clipping yields.

Growth and Morphological Characterization Study

Low- versus high-maintenance cultivars

Data taken during the course of the experiments showed the low- and high-maintenance cultivars consistently differed in rate of root growth and in clipping yields (Tables 8 and 9). The low-maintenance cultivars rooted to greater depths in the root profile and yielded greater clipping weights over all weeks in all experiments.

The data recorded at harvest showed that the low- and high-maintenance cultivars differed in many root and shoot growth characteristics (Tables 10 and 11). The low-maintenance cultivars produced greater total root mass, and greater root mass in the 0-20 and 20-40 cm sections of the root profile. The high-maintenance cultivars yielded a greater percentage of their total root weight in the 0-20 cm section, and the low-maintenance cultivars yielded a greater percentage of their total root weight in the 20-40 cm section of the root profile. Little root mass was produced in the 40-60 cm section of the root profile,

Table 6. Clipping yield data from the low-fertility, unirrigated cultivar field evaluation

	Sampling date-1987		
Cultivars	Sept. 1	Sept. 12	Oct. 15
	_____grams dry wt. m ⁻² _____		
	<u>Low-maintenance cultivars</u>		
S-21	23.0	12.3	7.7
S. D. Common	24.5	15.8	10.2
K3-162	27.3	17.5	7.2
Kenblue	28.3	17.7	8.8
Vantage	13.5	9.7	7.2
Mean	23.3	14.6	8.2
	<u>High-maintenance cultivars</u>		
I-13	1.1	2.8	1.3
Bonnieblue	11.5	5.7	3.8
A20	6.2	5.7	2.1
Columbia	5.2	3.9	2.4
Lovegreen	11.5	7.0	2.1
Mean	7.1	5.0	2.3
LSD (0.05)	14.8	9.2	4.8

Table 7. Mean squares from the analysis of clipping yield data from the low-fertility, unirrigated cultivar field evaluation

		Sampling date-1987		
Sources	df	Sept. 1	Sept. 12	Oct. 15
<hr/> grams dry wt. m ⁻² <hr/>				
Replication	2	322*	201	57**
Cultivars	9	292**	97*	32**
Contrast†	1	1974**	691**	260**
Among high‡	4	66	11	3
Among low§	4	95	40	5
Pooled error	18	74	29	8
Corrected total	29			

*, **Significant at the 5% or 1% level, respectively.

[†]Tests for differences between the combined means of the low- versus the high-maintenance cultivars.

[‡], [§]Tests for differences among the high-maintenance and among the low-maintenance cultivars, respectively.

Table 8. Root depth and clipping yield data from the low- and high-maintenance cultivar growth and morphological characterization study conducted in the greenhouse

Cultivars	Max. root depth								Clipping yields				
	Week								Week				
	3	4	5	6	7	8	9	10	5	6	7	8	9
	mm								mg				
	<u>Low-maintenance cultivars</u>												
S-21	120	206	252	286	324	350	376	424	31	50	63	122	96
S. D. Common	107	156	203	249	295	336	375	428	22	30	57	95	82
K3-162	99	154	199	252	298	322	370	423	32	48	72	123	98
Kenblue	102	161	196	232	264	304	348	390	33	45	68	103	88
Vantage	108	186	234	295	349	388	422	455	59	75	121	218	142
Mean	107	173	217	263	306	340	378	424	35	50	76	132	101
	<u>High-maintenance cultivars</u>												
I-13	61	82	110	134	172	200	236	307	0	2	3	18	17
Bonnieblue	105	155	194	237	258	288	314	374	3	11	18	56	65
A20	85	124	164	213	249	278	327	381	4	8	11	41	58
Columbia	84	118	162	182	215	239	282	333	6	10	17	60	56
Lovegreen	88	138	171	215	261	313	373	449	3	12	23	58	73
Mean	86	123	160	196	231	264	306	378	3	9	14	46	54
LSD (0.05)	17	35	40	44	51	55	59	56	16	20	29	49	36

Table 9. Means squares from the analysis of the root depth and clipping yield data from the cultivar growth and morphological characterization study conducted in the greenhouse

Source	df	Maximum root depth ($\times 10^3$)								Clipping yields ($\times 10^3$)				
		Week 3	Week 4	Week 5	Week 6	Week 7	Week 8	Week 9	Week 10	Week 5	Week 6	Week 7	Week 8	Week 9
		mm								mg				
Experiment	2	6.7**	48.3**	18.3**	14.8*	24.0*	70.6**	92.9**	174.1**	0.3	1.9	13.2*	28.2	27.9**
Rep(Exp.)	9	0.4	1.9	2.9	4.5	4.3	5.7	5.9	6.2	0.4	0.6	1.5	4.8	2.2
Cultivars	9	3.4**	14.8**	19.0**	27.0**	31.9**	35.5**	35.1**	28.6**	4.5**	7.1**	16.5**	39.4**	12.9*
Contrast†	1	15.3**	73.5**	96.8**	131.3**	168.5**	174.8**	154.5**	91.5**	31.0**	50.4**	113.5**	221.0**	66.7**
Among high‡	4	3.0**	8.9**	11.4*	19.2**	14.4*	23.7**	31.4**	35.0**	0.1	0.2	0.7	3.8	5.6
Among low§	4	0.8	5.9*	7.4	8.6	12.3	12.1	8.7	6.3	2.4*	3.2*	8.0	29.5*	6.8
Exp. * cultivar	18	0.4	1.8	2.8	3.4	4.5	4.7	6.0	7.0	0.6	0.8	3.4**	9.2**	3.9*
Pooled error	81	0.5	1.8	2.4	2.9	4.0	4.6	5.2	4.7	0.4	0.6	1.3	3.6	2.0
Corrected total	119													

*, **Significant at the 5% or 1% level, respectively.

†Tests for differences between the combined means of the low- versus the high-maintenance cultivars.

‡, §Tests for differences among the high-maintenance and among the low-maintenance cultivars, respectively.

Table 10. Root and shoot growth data from the low- and high-maintenance cultivar growth and morphological characterization study conducted in the greenhouse

Cultivars	Root wt. per section (cm)				Percent of total root wt. per section (cm)			Shoot weight		Lateral branch	Shoot to root ratio	Shoot dry to fresh wt. ratio		
	0-20	20-40	40-60	Total	0-20	20-40	40-60	Fresh	Dry					
	mg							mg					-no.-	mg/mg
	<u>Low-maintenance cultivars</u>													
S-21	172	55	3	230	75	24	1	1058	373	13	1.65	0.36		
S. D. Common	147	43	5	195	76	22	2	857	287	11	1.51	0.34		
K3-162	183	57	2	242	76	23	1	1062	361	15	1.52	0.35		
Kenblue	173	44	2	219	79	20	1	1048	354	15	1.61	0.35		
Vantage	209	75	12	296	71	26	3	1493	489	17	1.61	0.36		
Mean	177	55	5	238	75	23	2	1103	373	14	1.58	0.35		
	<u>High-maintenance cultivars</u>													
I-13	100	10	1	111	92	7	1	733	209	11	2.12	0.28		
Bonnieblue	114	25	2	141	82	16	2	1011	298	11	2.22	0.30		
A20	120	32	3	155	81	18	1	1005	319	13	2.13	0.32		
Columbia	148	26	1	175	87	13	0	1034	323	13	1.91	0.32		
Lovegreen	128	45	7	180	75	22	3	1202	361	15	2.17	0.30		
Mean	122	28	3	153	84	15	1	997	302	13	2.11	0.31		
LSD (0.05)	36	18	5	51	6	5	2	268	92	3	0.44	0.03		

Table 11. Mean squares from the analysis of the root and shoot growth data from the low- and high maintenance cultivar growth and morphological characterization study conducted in the greenhouse

Source	df	Percent of total											
		Root wt. per sections (cm)				root wt. per section (cm)			Shoot wt. (x 10 ³)		Lateral branch	Shoot to root ratio	Shoot dry to fresh wt. ratio
		0-20	20-40	40-60	Total	0-20	20-40	40-60	Fresh	Dry			
		mg							mg		no.-	mg/mg	
Experiment	2	31,333**	10,128**	478**	9,623	6,006**	4,305**	143.0**	3,145.7**	98.6*	320**	3.30**	0.090**
Rep(Exp.)	9	2,137	541	60*	3,319	65	51	6.6	151.9	12.5	10	0.45	0.001
Cultivars	9	14,253**	4,221**	127**	34,898**	507**	402**	14.5*	480.6	62.5	50	1.00*	0.009**
Contrast†	1	89,653**	22,032**	105	209,835**	2,076**	1,903**	3.8	NA†	NA	NA	8.40**	0.065**
Among high‡	4	3,700	2,303*	78*	9,306	550**	350**	16.5*	NA	NA	NA	0.18	0.004
Among low§	4	5,971	1,982*	183**	16,703	100	70	15.9*	NA	NA	NA	0.09	0.001
Exp. * cultivar	18	3,492*	657	19	6,607	81	68*	4.6	362.1**	27.6**	30**	0.30	0.002*
Pooled error	81	1,948	489	45	3,988	57	39	6.0	108.9	12.8	14	0.30	0.001
Corrected total	119												

*, **Significant at the 5% or 1% level, respectively.

†Tests for differences between the combined means of the low- versus the high-maintenance cultivars.

‡, §Tests for differences among the high-maintenance and among the low-maintenance cultivars, respectively.

‡NA indicates contrasts and tests among high and among low not performed where cultivar differences were nonsignificant.

and no differences were found between the low- and high-maintenance cultivars in root weight or percentage of the total root weight in this section of the profile. The low- and high-maintenance cultivars did not differ in shoot final fresh weight, shoot dry weight, and lateral branch number. The shoot to root ratio was calculated by dividing the final shoot dry weight by the total root dry weight, and low-maintenance cultivars were found to have a lower shoot to root ratio. The shoot dry weight was divided by the shoot fresh weight, and this ratio indicated that the shoot tissue of the low-maintenance cultivars contained less moisture.

Low- and high-maintenance cultivars also differed in all measured shoot morphological characteristics (Tables 12 and 13). The low-maintenance cultivars had narrower leaf blades, more leaf folding, longer leaf sheaths, a greater leaf angle from horizontal, and fewer leaves per shoot.

Variation among low- or high-maintenance cultivars

Data on total root weight, root weight in the 0-20 cm section of the rooting profile, clipping yields, shoot to root ratio, ratio of shoot dry weight to shoot fresh weight, and leaf folding separated clearly into low- and high-maintenance groups, and little variability was found among the cultivars within each group.

Generally, more variability was found among the high-maintenance cultivars than among the low-maintenance cultivars. Data on depth of rooting at all dates, root weight in the 20-40 cm section of the root profile, percentage of total root weight in the 0-20 and 20-40 cm sections

Table 12. Shoot morphology data from the low- and high-maintenance cultivar growth and morphological characterization study conducted in the greenhouse

Cultivars	Leaf			Sheath length	Leaves per shoot
	Width	Folding [†]	Angle [‡]		
	-mm-			-mm-	-no.-
<u>Low-maintenance cultivars</u>					
S-21	2.7	2.5	77.8	3.5	3.8
S. D. Common	2.4	2.8	76.2	3.2	3.9
K3-162	2.4	2.3	77.9	3.3	4.0
Kenblue	2.3	2.0	75.1	3.2	4.2
Vantage	3.0	2.7	73.5	3.4	3.9
Mean	2.5	2.5	76.1	3.3	4.0
<u>High-maintenance cultivars</u>					
I-13	3.0	4.3	69.2	2.1	4.3
Bonnieblue	3.0	3.8	63.3	2.7	4.9
A20	3.3	3.5	73.3	2.6	4.2
Columbia	3.0	3.8	64.8	2.7	4.5
Lovegreen	3.1	3.6	67.6	3.2	5.2
Mean	3.1	3.8	67.6	2.6	4.6
LSD (0.05)	0.2	0.5	4.0	0.3	0.4

[†]Leaf folding based on a scale of 1 to 5; 1 = tightly folded and 5 = flat leaf blades.

[‡]Leaf angle from horizontal measured from vertical youngest leaf to next youngest leaf.

Table 13. Mean squares from the analysis of the shoot morphology data from the low- and high-maintenance cultivar growth and morphological characterization study conducted in the greenhouse

Source	df	Leaf			Sheath length	Leaves per shoot
		Width	Folding [¶]	Angle [#]		
		—mm—			—mm—	—no.—
Experiment	2	0.04	2.36*	149*	0.98**	3.22**
Rep(Exp.)	9	0.06	0.95	18	0.46*	0.11
Cultivars	9	1.45**	7.19**	338**	2.24**	2.37**
Contrast [†]	1	8.22**	54.68**	2151**	13.20**	12.42**
Among high [‡]	4	0.19*	1.14	184**	1.62**	2.13**
Among low [§]	4	1.05**	1.26	42	0.31	0.30
Exp. * cultivar	18	0.06	0.53	26	0.16	0.22
Pooled error	81	0.04	0.42	25	0.16	0.20
Corrected total	119					

*, **Significant at the 5% or 1% level, respectively.

[†]Tests for differences between the combined means of the low- versus the high-maintenance cultivars.

[‡], [§]Tests for differences among the high-maintenance and among the low-maintenance cultivars, respectively.

[¶]Leaf folding based on a scale of 1 to 5; 1 = tightly folded and 5 = flat leaf blades.

[#]Leaf angle from horizontal measured from vertical youngest leaf to next youngest leaf.

of the root profile, leaf width, sheath length, leaf angle, and leaves per shoot also separated into distinct low- and high-maintenance groups. But, some variability within groups was observed.

The variability among the low-maintenance cultivars was largely due to the cultivar Vantage, and the variability among the high-maintenance cultivars was due primarily to the cultivars Lovegreen and I-13. Vantage produced more root weight in the 20-40 and 40-60 cm sections of the root profile and had wider leaf blades when compared to the other low-maintenance cultivars. I-13 had short leaf sheaths, and it had consistently low values for maximum root depth during the course of the experiments when compared to the other high-maintenance cultivars. I-13 produced little root mass in the lower sections of the root profile, and this resulted in a high percentage of the total root mass in the uppermost section of the profile relative to the other high-maintenance cultivars. Lovegreen had longer leaf sheaths, more leaves per tiller, and produced more root mass and a greater percentage of its total root mass in the 20-40 and 40-60 cm sections of the root profile than did the other high-maintenance cultivars.

Stomate Density Count Study

Low-maintenance cultivars had fewer stomata mm^{-2} on their abaxial leaf blade surface than the high-maintenance cultivars (Tables 14 and 15). There were no differences in the number of stomates mm^{-2} on the adaxial blade surface and in the ratio of the number of abaxial to adaxial stomata between low-and high-maintenance cultivars.

Table 14. Stomate density data from field grown low- and high-maintenance cultivars

Cultivars	Leaf blade surface		Ratio Abaxial/adaxial
	Abaxial	Adaxial	
	no. mm ⁻²		
	<u>Low-maintenance cultivars</u>		
S-21	109	240	0.46
S. D. Common	89	211	0.43
K3-162	83	242	0.34
Kenblue	102	250	0.41
Vantage	87	235	0.39
Mean	94	235	0.41
	<u>High-maintenance cultivars</u>		
I-13	145	266	0.55
Bonnieblue	91	208	0.43
A20	122	257	0.47
Columbia	136	256	0.53
Lovegreen	70	211	0.35
Mean	113	240	0.47
LSD (0.05)	26	40	0.09

Table 15. Mean squares from the analysis of the stomata density data from field grown low- and high-maintenance cultivars

Source	df	Leaf blade surface		Ratio Abaxial/adaxial
		Abaxial	Adaxial	
		no. mm ⁻²		
Experiment	1	1,198	1	0.02
Rep(Exp.)	6	549	600	0.01
Cultivar	9	4,704**	3,526	0.04
Contrast [†]	1	7,106*	NA [¶]	NA
Among high [‡]	4	8,684**	NA	NA
Among low [§]	4	1,984	NA	NA
Exp. * cultivar	9	820	2,626	0.02
Pooled error	54	667	1,615	0.01
Corrected total	79			

*, ** Significant at the 5% or 1% level, respectively.

[†] Tests for differences between the combined means of the low- versus the high-maintenance cultivars.

[‡], [§] Tests for differences among the high-maintenance and among the low-maintenance cultivars, respectively.

[¶] NA indicates contrast and test among high and among low not performed where cultivar differences were nonsignificant.

There were no differences among the low-maintenance cultivars, but there were differences among the high-maintenance cultivars in stomata number mm^{-2} on the abaxial leaf blade surface. This variability among the high-maintenance cultivars accounted for a large portion of the variability among all cultivars, and much of this variability was due to the high stomata number of I-13 and the low stomata number of Lovegreen.

Study on the Effect of N Level on Cultivar Shoot and Root Growth

Variation within the data of this study was mostly due to nitrogen (N) levels and cultivar differences (Tables 16,17, and 18). The N level had a substantial effect on shoot and root growth characteristics. Cultivars grown under low N rooted faster and produced greater root mass in the 20-40 cm section of the root profile, and had lower shoot to root ratios. Cultivars grown under high N produced greater clipping weights, shoot fresh and dry weights, and more lateral branches.

As was observed in the growth and morphological characterization study, the combined means of the low-maintenance and high-maintenance cultivars over both N levels differed in many root and shoot characteristics. Low-maintenance cultivars produced greater clipping yields, produced a greater percentage of their total root mass in the 20-40 section of the root profile, and had a lower shoot to root ratio than the high-maintenance cultivars. High-maintenance cultivars produced a greater percentage of their total root mass in the 0-20 cm section of the root profile.

Table 16. Root and shoot growth characteristics of two low- and two high-maintenance cultivars when grown under low and high nitrogen (N) levels

Parameter	Low-maintenance cultivars		High-maintenance cultivars	
	Low N	High N	Low N	High
Days to max. root depth	52	60	72	85
Root wt. per section, mg				
0-20 cm	91	89	98	76
20-40 cm	43	37	38	22
40-60 cm	25	20	16	11
Total root wt., mg	159	146	152	109
% total root wt. per section				
0-20 cm	58	61	65	68
20-40 cm	27	26	28	22
40-60 cm	15	13	10	11
Shoot fresh wt., mg	1052	2301	1209	2417
Shoot dry wt., mg	328	619	364	611
Lateral branch no.	15	24	14	25
Shoot to root ratio	2.10	4.31	2.71	5.74
Shoot dry to fresh wt ratio	0.31	0.27	0.30	0.25
Clipping wt., mg				
Week 4	30	59	15	28
Week 6	72	129	40	104
Week 8	63	130	36	98
Week 10	103	182	67	168

Table 17. Means squares from the analysis of the study on the effects of nitrogen (N) level on root depth and clipping yield characteristics of two low- and two high-maintenance cultivars

Source	df	Days to max. root depth	Clipping yield			
			Week 4	Week 6	Week 8	Week 10
			mg			
Experiment	1	5	1,701**	6,400*	4,200*	5,800**
Cultivars	3	2,895	3,700**	7,000**	6,400**	14,000**
Contrast	1	NA [†]	8,800**	12,700**	13,800**	10,300**
Error (a)	3	482	100	200	100	100
N	1	1,881*	7,400*	58,800**	66,300**	128,700**
N*cultivar	3	246	300	1,800	1,600	7,800
N*contrast	1	NA	800	200	200	1,900
Error (b)	4	181	500	2,200	1,300	1,300
Corrected total	63					

*, **Significant at the 5% and 1% level, respectively.

[†]NA indicates contrasts not performed when nonsignificant cultivar differences.

Table 18. Means squares from the analysis of the study on the effect of nitrogen (N) level on root and shoot growth characteristics of two low- and two high-maintenance cultivars

		Percent of total root wt.												
		Root wt. per section (cm)				per section (cm)			Shoot wt. (x 10 ³)		Lateral	Shoot to	Shoot dry to	
Source	df	0-20	20-40	40-60	Total	0-20	20-40	40-60	Fresh	Dry	branch	root	fresh wt.	
		mg							mg		no.-	mg/mg		
Experiment	1	1,712*	2,849*	2,316*	20,413*	1049*	159*	395*	7,501**	574**	124**	44.90	0.0001	
Cultivars	3	8,168**	910	657	17,129*	442*	159*	104	1,033*	70	62*	7.81*	0.0010	
Contrast	1	186	NA	NA	8,213	798*	191*	NA	298	NA	1	16.40*	NA	
Error (a)	3	98	146	139	1,039	38	7	15	110	12	7	0.65	0.0003	
N	1	2,316	1,969*	276	11,908	119	117	1	24,145**	1,155**	1,550**	110.26**	0.0356**	
N*cultivar	3	732	519	140	2,957	217	69	61	182	14	8	1.22	0.0002	
N*contrast	1	NA†	NA	NA	NA	5	20	NA	NA	NA	NA	0.02	NA	
Error (b)	4	1,219	114	52	2,082	49	57	18	685	52	23	1.58	0.0001	
Corrected total	63													

*, **Significant at the 5% and 1% level, respectively.

†NA indicates contrasts not performed where cultivar differences were nonsignificant.

Little variation occurred within the data of this study due to the N * cultivar interaction, and no data exhibited a significant N * low-maintenance versus high-maintenance cultivar interaction. This indicated that the differences between the low-maintenance and high-maintenance cultivars did not change when grown under the two N levels.

DISCUSSION

Plant characteristics identified as drought tolerance adaptations include root characteristics that increase absorption, shoot characteristics that decrease transpiration, and low shoot to root ratios (29,32). Deep, extensive root systems are the principle method by which monocots increase absorption (32). Shoot characteristics that reduce transpiration include decreased shoot surface area, increased shoot canopy resistance to evapotranspiration, and few, small, and responsive stomata (29). Shoot characteristics of grasses that function to decrease shoot surface area include leaf folding or rolling (32), few leaves per unit area, narrow leaf blade width, low shoot density, and slow leaf elongation rate (28). Shoot characteristics of grasses that function to increase shoot canopy resistance to evapotranspiration include compact growth habit (32), horizontal leaf orientation, and high shoot density (28).

Kentucky bluegrass cultivars have been shown to differ in many shoot (36,41) and root growth characteristics (5). Observations from this investigation have shown that the differences in many of these characteristics separate clearly into low- and high-maintenance groups. These differences appear to be real and measurable, and observations and data from the field and the nutrient solution greenhouse study show the differences appear to be consistent over high and low nitrogen fertility levels.

Data from this investigation showed low-maintenance Kentucky bluegrass cultivars rooted faster, and they produced greater root mass and

a greater percentage of their total root mass lower in the root profile. Low-maintenance cultivars maintained a lower shoot to root ratio. With the exception of their rapid rate of leaf elongation, low-maintenance cultivars possessed shoot characteristics that reduced their shoot surface area. High-maintenance cultivars possessed shoot characteristics that increased their canopy resistance to evapotranspiration.

Differences in Kentucky bluegrass cultivar growth and morphology on the whole plant level may play a more important role in reducing transpiration loss than the differences on the cellular level, such as stomatal density (13,24). This study showed low-maintenance cultivars had fewer stomata on their abaxial leaf blade surface, but much of the difference among cultivars in abaxial stomata density was explained by the variation among cultivars from the high-maintenance group.

Variation did occur both among the low-maintenance and high-maintenance cultivars in some of the data collected during this investigation. But, considering the diverse origin and nature of Kentucky bluegrass cultivars, some variation among cultivars within each maintenance group was to be expected. The high-maintenance cultivar Lovegreen was the only cultivar that displayed characteristics of both maintenance groups.

It is probably a combination of many adaptations that allow some Kentucky bluegrass cultivars to out perform others when grown under low fertility, unirrigated culture. The shoot characteristics known to be important drought stress tolerance mechanisms were not common only to the low-maintenance cultivars in this study. The root production, root distribution, low shoot to root ratio, and low shoot moisture content

characteristics were consistently found among the low-maintenance cultivars, and these adaptations may best explain the reliable performance of these cultivars under low fertility, unirrigated management.

The ultimate goal of cultivar development programs is to select and develop cultivars with desired shoot characteristics that are well adapted to a wide range of management conditions and environmental stresses. The identification of characteristics common to the low-maintenance cultivars that have no effect on aesthetics, such as the rooting characteristics and shoot to root ratios, may be beneficial for the selection and development of improved Kentucky bluegrass cultivars, and also for the identification of existing improved Kentucky bluegrass cultivars more tolerant to a wide range of management practices and environmental conditions.

REFERENCES

1. Arcioni, S., M. Falcinelli, and D. Mariotti. 1985. Ecological adaptation in Lolium perenne L.: Physiological relationships among persistence, carbohydrate reserves and water availability. Can. J. Plant Sci. 65:615-624.
2. Aronson, L. J., A. J. Gold, and R. J. Hull. 1987. Cool-season turfgrass responses to drought stress. Crop Sci. 27:1261-1266.
3. Baker, B. J. and G. A. Jung. 1968. Effect of environmental conditions on the growth of four perennial grasses. Agron. J. 60:155-158.
4. Beard, J. B. 1973. Turfgrass: Science and Culture. Prentice-Hall, Inc., Englewood Cliffs, New Jersey.
5. Boeker, P. 1974. Root development of selected turfgrass species and cultivars. pp. 55-61. In E. C. Roberts (ed.) Proceedings of the 2nd Int. Turfgrass Research Conference. American Society of Agronomy and Crop Science Society of America, Madison, WI.
6. Burton, G. W., G. M. Prine, and J. E. Jackson. 1957. Studies of drought tolerance and water use of several southern grasses. Agron. J. 49:498-503.
7. Campbell, G. S., R. I. Papendick, E. Rabie, and A. J. Shayo-Ngowi. 1979. A comparison of osmotic potential, elastic modulus, and apoplastic water in leaves of dryland winter wheat. Agron. J. 71:31-36.
8. Carrier, L. and K. S. Bort. 1916. The history of Kentucky bluegrass and white clover in the United States. J. Am. Soc. Agron. 8:256-266.
9. Carroll, J. C. 1943. Effects of drought, temperature, and nitrogen on turfgrasses. Plant Physiol. 18:19-36.
10. Daniel, W. H. and R. P. Freeborg. 1979. Turf manager's handbook. Harvest Publishing Co., Cleveland, Ohio.
11. Derera, N. F., D. R. Marshall, and L. N. Balaam. 1969. Genetic variability in root development in relation to drought tolerance in spring wheats. Exp. Agric. 5:327-337.
12. Dernoeden, P. H. and J. D. Butler. 1978. Drought resistance of Kentucky bluegrass cultivars. HortSci. 13(6):667-668.

13. Dernoeden, P. H. and J. D. Butler. 1979. Relation of various plant characters to drought resistance of Kentucky bluegrass. *HortSci.* 14(4):511-512.
14. Fereres, E., E. Acevedo, P. W. Henderson, and T. C. Hsiao. 1978. Seasonal changes in water potential and turgor maintenance in sorghum and maize under water stress. *Physiol. Plant.* 44:261-267.
15. Fergus, E. N. and R. C. Buckner. 1973. The bluegrasses and redtop. p. 243-253. In M. E. Heath, D. S. Metcalfe, and R. F. Barnes (eds.) *Forages*. Iowa State University Press, Ames, Iowa.
16. Funk, C. R. and R. E. Engel. 1984. Kentucky bluegrasses and their culture. *Weeds, Trees, and Turf* 23(6):46-62.
17. Gaskin, T. A. 1966. Testing for drought and heat resistance in Kentucky bluegrass. *Agron. J.* 58:461-462.
18. Hanson, A. A., F. V. Juska, and G. W. Burton. 1969. Species and varieties. p. 370-409. In A. A. Hanson and F. V. Juska (eds.) *Turfgrass science*. American Society of Agronomy, Inc., Madison, Wisconsin.
19. Hurd, E. A. 1968. Growth of roots of seven varieties of spring wheat at high and low moisture levels. *Agron. J.* 60:201-205.
20. Iowa State University Cooperative Extension Service. 1984. 1984 Iowa turfgrass research report FG-450.
21. Iowa State University Cooperative Extension Service. 1985. 1985 Iowa turfgrass research report FG-451.
22. Iowa State University Cooperative Extension Service. 1986. 1986 Iowa turfgrass research report FG-452.
23. Iowa State University Cooperative Extension Service. 1987. 1987 Iowa turfgrass research report FG-453.
24. Johns, D., Jr., J. B. Beard, and C. H. M. van Bavel. 1983. Resistance to evapotranspiration from a turfgrass canopy. *Agron J.* 75:419-422.
25. Jones, M. M. and N. C. Turner. 1978. Osmotic adjustment in leaves of sorghum in response to water deficits. *Plant Physiol.* 61:122-126.
26. Jordan, W. R. and F. R. Miller. 1980. Genetic variability in sorghum root systems: Implications for drought tolerance. In N. C. Turner and P. J. Kramer (eds.) *Adaptation of plants to water and high temperature stress*. John Wiley and Sons, Inc., New York, New York.

27. Julander, O. 1945. Drought resistance in range and pasture grasses. *Plant Physiol.* 20:573-599.
28. Kim, K. S. and J. B. Beard. 1988. Comparative turfgrass evapotranspiration rates and associated plant morphological characteristics. *Crop Sci.* 28:328-331.
29. Kramer, P. J. 1980. Drought, stress, and the origin of adaptations. pp. 7-20. In N. C. Turner and P. J. Kramer (eds.) *Adaption of plants to water and high temperature stress.* John Wiley and Sons, Inc., New York, New York.
30. Laude, H. M. 1953. The nature of summer dormancy in perennial grasses. *Bot. Gaz.* 114(3):284-292.
31. Lehman, V. G. and M. C. Engelke. 1985. A rapid screening technique for genetic variability in turfgrass root systems. pp. 770-776. In F. Lemaire (ed.) *Proceedings of the 5th Int. Turfgrass Conference.* INRA Publications and The International Turfgrass Society, Versailles, France.
32. Levitt, J. 1980. Responses of plants to environmental stresses. Volume II: Water, radiation, salt, and other stresses. Academic Press, New York, New York.
33. May, L. H. and F. L. Milthorpe. 1962. Drought resistance of crop plants. *Field Crop Abstr.* 15(3):171-179.
34. Minner, D. D. and J. D. Butler. 1985. Drought tolerance of cool season turfgrasses. pp. 199-212. In F. Lemaire (ed.) *Proceeding of the 5th Int. Turfgrass Research Conference.* INRA Publications and the International Turfgrass Society, Versailles, France.
35. Minner, D. D., P. H. Dernoden, D. J. Wehner, and M. S. McIntosh. 1983. Heat tolerance screening of field-grown cultivars of Kentucky bluegrass and perennial ryegrass. *Agron. J.* 75:772-775.
36. Nittler, L. W. 1966. Seedling characteristics useful in identifying Kentucky bluegrass. *Farm Res.* 32(1):14-15.
37. Nus, J. L. and C. F. Hodges. 1985. Differential sensitivities of various parts of Kentucky bluegrass to water stress. *HortSci.* 21(4):1014-1015.
38. Ofir, M. and D. Kerem. 1982. The effects of temperature and photoperiod on the onset of summer dormancy in Poa bulbosa L. *Ann. Bot.* 50:259-264.
39. Pellett, H. M. and E. C. Roberts. 1963. Effects of mineral nutrition on high temperature induced growth of Kentucky bluegrass. *Agron. J.* 55:473-476.

40. Scarth, G. W., H. B. Gooding, and M. Shaw. 1947. Factors influencing growth and summer dormancy in Taraxacum kok-saghyz. Can. J. Res. 25:27-42.
41. Schery, R. W. 1965. This remarkable Kentucky bluegrass. Ann. Mo. Bot. Gard. 52:441-451.
42. Shearman, R. C. and J. B. Beard. 1972. Stomatal density and distribution in Agrostis as influenced by species, cultivar, and leaf blade surface and position. Crop Sci. 12:822-823.
43. Sheffer, K. M. 1979. Responses of three cool-season turfgrass species to heat and moisture stress. Ph.D. Dissertation. Univ. of Missouri, Columbia.
44. Taylor, L. H., J. F. Shoulders, A. J. Powell, Jr., and R. E. Schmidt. 1974. Development of improved Kentucky bluegrass cultivars for the transition zone of the eastern United States. pp. 89-92. In E. C. Roberts (ed.) Proceedings of the 2nd Int. Turfgrass Research Conference. American Society of Agronomy and Crop Science Society of America, Madison, WI.
45. Turgeon, A. J. 1985. Turfgrass management. Prentice-Hall, Inc., Reston, Virginia.
46. U. S. Department of Agriculture Agricultural Research Service and the Maryland State Turfgrass Council. 1986. 1980 National Kentucky bluegrass test final report PGGI No. 86-16.
47. Vegis, A. 1964. Dormancy in higher plants. Ann. Rev. Plant Physiol. 15:185-224.
48. Wallner, S. J., M. R. Becwar, and J. D. Butler. 1982. Measurement of turfgrass heat tolerance in vitro. J. Am. Soc. Hort. Sci. 107(4):608-613.
49. Watschke, T. L. and R. E. Schmidt. 1969. Performance of some Kentucky bluegrasses under high temperatures. pp. 258-259. In Sports Turf Research Institute (eds.) Proceedings of the 1st Int. Turfgrass Research Conference. Sports Turf Research Institute and the International Turfgrass Society, Bingley, Yorkshire, U. K.
50. Watschke, T. L., R. E. Schmidt, E. W. Carson, and R. E. Blaser. 1972. Some metabolic phenomena of Kentucky bluegrass. Crop Sci. 12:87-90.

51. Wehner, D. J. and T. L. Watschke. 1981. Heat tolerance of Kentucky bluegrasses, perennial ryegrasses, and annual bluegrass. *Agron. J.* 73:79-84.
52. Whalley, R. D. B. and A. A. Davidson. 1968. Physiological aspects of drought dormancy in grasses. *Proc. Ecol. Soc. Aust.* 3:17-19.

ACKNOWLEDGMENTS

My sincerest appreciation goes to Dr. Nick Christians for his sound guidance and support in making my advanced education possible. Nick is a valuable asset to I.S.U. and to the entire turf industry. I also would like to thank my graduate committee members Drs. Mike Agnew, Clint Hodges, and Walt Wedin for their support and advice in preparing and editing this manuscript. I am grateful to Dr. Paul Hinz for his statistical advice, and to Glenn Pearston for his computer expertise. Appreciation is expressed to Dr. Charles Hall and the Horticulture department faculty and staff for their assistance throughout my program of study. I am also thankful for the good humor and moral support of my officemates, especially my great friends Zac Reicher, Diane Shogren, and Gary Polking. Ultimately, I want to thank my wife, Diane, my daughter, Bailey, and my parents and family for their continuing love, patience, and support.