


Graphical Abstract

To create your abstract, type over the instructions in the template box below.
Fonts or abstract dimensions should not be changed or altered.

Acyl furans from cyclohexane-1,3-diones – A synthesis of hibiscone C

Yang Qu, George A Kraus*

Leave this area blank for abstract info.





Acyl furans from cyclohexane-1,3-diones – A synthesis of hibiscone C

Yang Qu, George A. Kraus*

Department of Chemistry, Iowa State University, Ames, IA, 50010

ARTICLE INFO

Article history:

Received

Received in revised form

Accepted

Available online

Keywords:

Acyl furan

Cyclohexane-1,3-dione

Hibiscone C

Dithiane

Synthesis

ABSTRACT

A direct route to acyl furans was developed using dithiane anion addition followed by deprotection/aromatization. This led to an efficient synthesis of hibiscone C.

© 2009 Elsevier Ltd. All rights reserved.

A number of natural products contain the acyl furan subunit, including viridin (**1**), halenaquinone (**2**), hibiscone C (**3**) and hibiscone D (**4**), shown in Figure 1. There have been total syntheses¹ of **1** and a number of innovative approaches² to the ring structure of **1**. Halenaquinone is a novel RAD51 inhibitor that specifically inhibits the RAD51-dsDNA binding.³ Several researchers have reported total syntheses⁴ of **2** and approaches⁵ to **2**. Hibiscone C competitively inhibits phosphatidylinositol-3-kinase activity in intact cells.⁶ Since the classic synthesis by Smith in 1982, it has been synthesized by Kraus, by Goess and by Lu.⁷ Hibiscone D inhibits superoxide anion generation by human neutrophils.⁸





Figure 1. Structures of acyl furans


Our approach began with enone **6**, as shown in the retrosynthetic analysis in Scheme 1. Enones such as **6** are readily constructed from cyclohexane-1,3-diones. Initially, we used

enone **6** (R = H)⁹ to identify optimal conditions. Interestingly, **5** (R = H) has not previously been reported.



Scheme 1. Retrosynthetic analysis


As shown in Scheme 2, the synthesis of furan **5** from **6** required a one-carbon nucleophile. The dithiane anion reacted efficiently with **6** to provide the alkoxide which was quenched with 2M aqueous HCl to afford **7** in 68% yield. The dithiane alcohol **7** was oxidized with bis(trifluoroacetoxy)iodobenzene in wet methanol to provide an acetal **8** which could be converted into **5** using PTSA in 75% yield over two steps.



Scheme 2. Synthesis of core structure **5**

* Corresponding author. Tel.: +0-000-000-0000; fax: +0-000-000-0000; e-mail: gakraus@iastate.edu

With a good route to furan **5**, we focused on the pathway to hisbicone C shown in Scheme 3. Commercially available 5-isopropyl-1,3-cyclohexanedione was converted into an acetal which was alkylated with allyl bromide to give **9** in 71% yield. The coupling constant of 4.8 Hz confirmed that the two groups were trans. Reaction with dithianyl lithium followed by quenching with HCl afforded enone **10** in 68% yield. Dithiane removal and aromatization furnished **11** in 75% yield. Conversion of the alkene to the primary alcohol using common hydroboration-oxidation conditions gave low yields. Fortunately, reaction with catechol borane mediated by Wilkenson's catalyst¹⁰ gave a 55% yield of compound **12**. Oxidation of alcohol **12** with excess PCC generated an aldehyde which cyclized to a furanyl alcohol which was oxidized in situ to afford diacylfuran **13**. Since **12** was an intermediate in the synthesis of hisbicone C by Lu⁷, this represents a formal total synthesis of **3**.



Scheme 3. Synthesis of compound **13**

The successful synthesis of hisbicone C showcases a direct route to 2,4-diacyl furans. The route is scalable and the individual reactions are operationally convenient.

Acknowledgments

We thank the Iowa State Department of Chemistry for partial support for YQ.

References and notes

- (a) Bel, M. D.; Abela, A. R.; Ng, J. D.; Guerrero, C. A. *J. Am. Chem. Soc.* **2017**, *139*(20), 6819. (b) Mizutani, T.; Honzawa, S.; Tosaki, S.-Y.; Shibasaki, M. *Angew. Chem.* **2002**, *114*(24), 4874. (c) Shigehisa, H.; Mizutani, T.; Tosaki, S.-Y.; Ohshima, T.; Shibasaki, M. *Tetrahedron* **2005**, *61*(21), 5057. (d) Anderson, E. A.; Alexanian, E. J.; Sorensen, E. J. *Angewandte Chemie International Edition* **2004**, *43*(15), 1998.
- (a) Jacobi, P. A.; Mascall, K. C. *Heterocycles* **2014**, *88*(2), 1527. (b) Jacobi, P. A.; Könekamp, T.; Mascall, K. C.; O'Connor, R. T.; Onyango, E. O.; Sessions, E. H. *Advances in Heterocyclic Chemistry* **2013**, *119*. (c) Onyango, E. O.; Jacobi, P. A. *J. Org. Chem.* **2012**, *77*(17), 7411. (d) Keay, B.; Muller, K. *Synlett* **2008**, *2008*(8), 1236. (e) Sessions, E. H.; Jacobi, P. A. *Org. Lett.* **2006**, *8*(18), 4125. (f) Souza, F. E. S.; Rodrigo, R. *Chem. Comm.* **1999**, *19*, 1947.
- Takaku, M.; Kainuma, T.; Ishida-Takaku, T.; Ishigami, S.; Suzuki, H.; Tashiro, S.; Soest, R. W. M. V.; Nakao, Y.; Kurumizaka, H. *Genes to Cells* **2011**, *16*(4), 427.
- (a) Goswami, S.; Harada, K.; El-Mansy, M. F.; Lingampally, R.; Carter, R. G. *Angew. Chem.* **2018**, *130*(29), 9255. (b) Kienzler, M. A.; Suseno, S.; Trauner, D. *J. Am. Chem. Soc.* **2008**, *130* (27), 8604. (c) Harada, N.; Sugioka, T.; Ando, Y.; Uda, H.; Kuriki, T. *J. Am. Chem. Soc.* **1988**, *110*(25), 8483. (d) Kojima, A.; Takemoto, T.; Sodeoka, M.; Shibasaki, M. *J. Org. Chem.* **1996**, *61*(15), 4876. (e) Kojima, A.; Takemoto, T.; Sodeoka, M.; Shibasaki, M. *Synthesis* **1998**, 581–589. (f) Souza, F. E. S.; Rodrigo, R. G. A. *J. Org. Chem.* **2001**, *66*(12), 4462.
- (a) Cristofoli, W. A.; Keay, B. A. *Synlett* **1994**, *1994*(08), 625. (b) Harada, N.; Sugioka, T.; Uda, H.; Kuriki, T.; Kobayashi, M.; Kitagawa, I. *J. Org. Chem.* **1994**, *59*(22), 6606. (c) Toyooka, N.; Nagaoka, M.; Kakuda, H.; Nemoto, H. *Synlett* **2001**, *2001*(07), 1123. (d) Toyooka, N.; Nagaoka, M.; Sasaki, E.; Qin, H.; Kakuda, H.; Nemoto, H. *Tetrahedron* **2002**, *58*(30), 6097. (e) Wakefield, B.; Halter, R. J.; Wipf, P. *Org. Lett.* **2007**, *9*(16), 3121.
- Besley, C.; Rhinehart, D. P.; Ammons, T.; Goess, B. C.; Rawlings, J. S. *Bio. Med. Chem. Lett.* **2017**, *27*(14), 3087.
- (a) Koft, E. R.; Smith, A. B. *J. Am. Chem. Soc.* **1982**, *104*, 5568–5570. (b) Koft, E. R.; Smith, A. B. *J. Am. Chem. Soc.* **1984**, *106*, 2115–2121. (c) Kraus, G. A.; Wan, Z. *Synlett* **2000**, *2000*(03), 363. (d) Ungureanu, S.; Meadows, M.; Smith, J.; Duff, D. B.; Burgess, J. M.; Goess, B. C. *Tetrahedron Letters* **2011**, *52*(13), 1509. (e) Lu, Y.; Yuan, H.; Zhou, S.; Luo, T. *Org. Lett.* **2017**, *19*(3), 620.
- Cheng, L.; Liao, H.; Chen, L.; Wang, S.; Kuo, Y.; Chung, M.; Chen, J. *Fitoterapia* **2017**, *117*, 16–21.
- Parida, A.; Sharique, M.; Kakde, B. N.; Ghosh, S.; Bisai, A. *Synthesis* **2015**, *47*(19), 2965–2970.
- Delorbe, J. E.; Lotz, M. D.; Martin, S. F. *Org. Lett.* **2010**, *12*(7), 1576.
- Experimental procedure for the synthesis of **3-(1,3-dithian-2-yl)-2-(hydroxymethyl)cyclohex-2-en-1-one (7)**: 1,3-dithiane (574 mg, 4.77 mmol) was dissolved in 15 mL freshly distilled THF in a round bottom flask and cooled to -78°C . n-BuLi solution (2.5M, 4.4 mmol) was added dropwise and the mixture was stirred at -78°C for 1.5 hours. **6** (566 mg, 3.67 mmol) in 10 mL THF was added dropwise to the flask at -78°C , warm up to room temperature and let it stir for additional 6 hours. Quench the reaction with 2M aqueous HCl (5 mL) and stir for overnight. Extract with Ethyl Acetate and dried over sodium sulfate. Concentration and flash column chromatography gave compound **7** in 72% yield; $^1\text{H NMR}$ (400 MHz, Chloroform-*d*) δ = 5.31 (s, 1H), 4.47 (d, $J=5.9$, 2H), 3.09–2.97 (m, 2H), 2.88 (ddd, $J=14.6$, 4.3, 3.0, 2H), 2.63 (t, $J=6.0$, 3H), 2.42 (dd, $J=7.5$, 6.0, 2H), 2.15 (ddt, $J=14.2$, 4.3, 2.0, 1H), 2.03–1.93 (m, 2H); $^{13}\text{C NMR}$ (100 MHz, Chloroform-*d*) δ = 156.2, 56.4, 50.6, 37.5, 31.0, 28.0, 25.0, 22.1; HRMS (ESI-QTOF) calcd for $\text{C}_{11}\text{H}_{16}\text{O}_2\text{S}_2$ [$\text{M}+\text{H}$] $^+$ 245.0664, found 245.0663.
- Experimental procedure for the synthesis of **6,7-dihydroisobenzofuran-4(5H)-one (5)**: **7** (319 mg, 1.31 mmol) was dissolved in methanol and water mixture solvent (4.5 mL methanol, 0.5 mL water). To the mixture was added bis(trifluoroacetoxy)iodobenzene (1.127 g, 2.62 mmol) portionwise at 0°C , warm up to room temperature and let it stir for 1 hour. P-TSA (500mg, 2.62 mmol) was added and the mixture was stirred at 60°C for 6 hours. The reaction was cooled to room temperature before water was added, extract with Ethyl Acetate and dried over sodium sulfate. Concentration and flash column chromatography gave compound **5** in 82% yield; $^1\text{H NMR}$ (400 MHz, Chloroform-*d*) δ = 7.95 (d, $J=1.4$, 1H), 7.25 (q, $J=1.4$, 1H), 2.66 (td, $J=6.2$, 1.4, 3H), 2.52–2.42 (m, 2H), 2.08–1.96 (m, 2H); $^{13}\text{C NMR}$ (100 MHz, Chloroform-*d*) δ = 195.2, 144.2, 138.1, 124.6, 124.0, 39.7, 24.0, 19.6; HRMS (ESI-QTOF) calcd for $\text{C}_8\text{H}_8\text{O}_2$ [$\text{M}+\text{H}$] $^+$ 137.0597, found 137.0601.
- 4-allyl-3-(1,3-dithian-2-yl)-2-(hydroxymethyl)-5-isopropylcyclohex-2-en-1-one (10)**: compound **10** was synthesized in 68% yield using the procedure for compound **7**; $^1\text{H NMR}$ (400 MHz, Chloroform-*d*) δ = 5.80 (dddd, $J=17.0$, 10.2, 8.5, 5.3, 1H), 5.18–5.08 (m, 2H), 4.89 (s, 1H), 4.65 (d, $J=2.8$, 2H), 3.04–2.93 (m, 2H), 2.90–2.82 (m, 2H), 2.81–2.56 (m, 3H), 2.49–2.42 (m, 1H), 2.30–2.21 (m, 1H), 2.15 (dt, $J=13.7$, 4.5, 2.2, 1H), 1.96–1.85 (m, 1H), 1.73–1.65 (m, 1H), 1.62–1.53 (m, 1H); $^{13}\text{C NMR}$ (100 MHz, Chloroform-*d*) δ = 200.2, 156.1, 137.6, 136.0, 117.4, 57.1, 51.7, 42.2, 41.8, 36.6, 35.9, 32.4, 32.0, 29.2, 25.1, 21.3, 20.8; HRMS (ESI-QTOF) calcd for $\text{C}_{17}\text{H}_{26}\text{O}_2\text{S}_2$ [$\text{M}+\text{H}$] $^+$ 327.1447, found 327.1445.
- 7-allyl-6-isopropyl-6,7-dihydroisobenzofuran-4(5H)-one (11)**: compound **11** was synthesized in 75% yield using the procedure for compound **5**; $^1\text{H NMR}$ (400 MHz, Chloroform-*d*) δ = 7.95 (d, $J=1.4$, 1H), 7.30 (t, $J=1.4$, 1H), 5.79 (dddd, $J=17.4$, 9.7, 7.5, 6.4, 1H), 5.10 (dq, $J=4.9$, 1.7, 1H), 5.07 (d, $J=2.0$, 1H), 2.91 (d, $J=5.9$, 1H), 2.63–2.47 (m, 2H), 2.43–2.36 (m, 2H), 1.86–1.77 (m,

2H); ^{13}C NMR (100 MHz, Chloroform- d) δ = 194.9, 143.5, 139.0, 135.3, 127.3, 123.5, 117.1, 45.3, 38.6, 37.1, 33.0, 27.7, 21.1, 17.9; HRMS (ESI-QTOF) calcd for $\text{C}_{14}\text{H}_{18}\text{O}_2$ $[\text{M}+\text{H}]^+$ 219.1380, found 219.1378.

15. **5-isopropyl-5,5a,6,7-tetrahydro-3H-naphtho[1,8-bc]furan-3,8(4H)-dione (13):** **12** (115 mg, 0.49 mmol) was dissolved in 2 mL DCM, to the solution PCC (431 mg, 1.96 mmol) was added and the mixture was stirred for 3 hours at room temperature. Short column was performed before drying over sodium sulfate. Concentration and Flash column chromatography gave compound **13** in 65% yield; ^1H NMR (400 MHz, CDCl_3) δ = 8.09 (s, 1H), 2.96 (td, J = 11.4, 4.8 Hz, 1H), 2.72 – 2.65 (m, 2H), 2.60 (dd, J = 16.8, 2.8 Hz, 1H), 2.45 (dq, J = 12.5, 4.1 Hz, 1H), 2.34 (dd, J =

16.8, 13.3 Hz, 1H), 2.12 – 2.03 (m, 1H), 1.90 (ddd, J = 13.7, 5.5, 2.7 Hz, 1H), 1.75 (ddd, J = 24.4, 12.5, 5.6 Hz, 1H), 1.01 (d, J = 7.0 Hz, 3H), 0.96 (d, J = 6.9 Hz, 3H); ^{13}C NMR (101 MHz, Chloroform- d) δ = 193.4, 185.0, 147.4, 145.8, 145.6, 48.0, 39.9, 38.7, 34.2, 29.7, 26.7, 20.9, 15.5; HRMS (ESI-QTOF) calcd for $\text{C}_{14}\text{H}_{16}\text{O}_3$ $[\text{M}+\text{H}]^+$ 233.1172, found 233.1173.

[Click here to remove instruction text...](#)