



# Role of dispersal in resistance evolution and spread

Nicholas J Miller<sup>1</sup> and Thomas W Sappington<sup>2</sup>

Gene flow via immigration affects rate of evolution of resistance to a pest management tactic, while emigration from a resistant population can spread resistance alleles spatially. Whether resistance detected across the landscape reflects ongoing *de novo* evolution in different hotspots or spread from a single focal population can determine the most effective mitigation strategy. Pest dispersal dynamics determine the spatio-temporal scale at which mitigation tactics must be applied to contain or reverse resistance in an area. Independent evolution of resistance in different populations appears common but not universal. Conversely, spatial spread appears to be almost inevitable. However, rate and scale of spread depends largely on dispersal dynamics and interplay with factors such as fitness costs, spatially variable selection pressure and whether resistance alleles are spreading through an established population or being carried by populations colonizing new territory.

## Addresses

<sup>1</sup> Department of Biology, Illinois Institute of Technology, 298 Life Science Building, 3101 S. Dearborn St., Chicago, IL 60616, USA

<sup>2</sup> USDA, Agricultural Research Service, Corn Insects and Crop Genetics Research Unit, Genetics Laboratory, Iowa State University, Ames, IA 50011, USA

Corresponding author: Sappington, Thomas W  
([Tom.Sappington@ars.usda.gov](mailto:Tom.Sappington@ars.usda.gov))

Current Opinion in Insect Science 2017, 21:68–74

This review comes from a themed issue on **Pests and resistance**

Edited by **Thomas W Sappington** and **Nicholas John Miller**

For a complete overview see the [Issue](#) and the [Editorial](#)

Available online 22nd May 2017

<http://dx.doi.org/10.1016/j.cois.2017.04.005>

2214-5745/Published by Elsevier Inc.

## Introduction

Insect dispersal plays a pivotal role in both the evolution of resistance to an insecticide or other management tactic at a location, and in the rate and pattern of its spatial spread [1,2]. Dispersal by individual insects is the fundamental process by which resistance genes move across a landscape. Consider a population where resistant individuals are favored by local selection. At its simplest level, immigration of susceptible individuals decreases the frequency of resistance alleles in the receiving population and thus slows the local rate of resistance evolution. Conversely, immigration of resistant individuals increases

resistance allele frequency and thus increases the rate of resistance evolution in that population. Immigration in one population presupposes emigration from another, and the rate of emigration of individuals from a resistant source population affects the rate of resistance spread in the landscape (Figure 1). The relative roles of *de novo* resistance evolution vs. subsequent spread can be of great practical importance because it determines the most appropriate mitigation strategy to be employed. The spatial scale at which mitigation tactics must be applied within the larger landscape to contain or reverse resistance that has evolved in a local ‘hotspot’ depends on dispersal dynamics of the species. As critical as these rates of inflow and outflow are to all aspects of insect resistance management (IRM), they are poorly understood for most species. Accordingly, the complex dynamics of resistance evolution and spread are likewise difficult to characterize.


A distinction can be made between dispersal that moves resistance genes between existing populations by gene flow, and dispersal that moves resistance genes into new territory as the insect colonizes previously unoccupied habitat. The first involves a process that can be likened to an invasion of resistance alleles into parts of the larger metapopulation that originally had none. The second, colonization of new habitat by insects carrying resistance alleles conflates the spatial invasion of alleles via gene flow with the genetic consequences associated with geographic invasion by a species, such as bottlenecks. The evolutionary ecological outcomes of both processes can be quite interesting.

## Spread of resistance among existing populations

### Case study: *Culex pipiens* resistance allele clines

An especially well studied case of the spatial and temporal distribution of resistance alleles in established populations comes from the mosquito *Culex pipiens* in the south of France [3–6]. In this case, the dynamics of resistance were studied at a small geographical scale relative to dispersal distances. Consequently, alleles have usually been observed at migration-selection equilibrium. In this region there is an annual program of mosquito control whereby breeding sites within approximately 20 km of the Mediterranean coast are treated with insecticides every breeding season. This program relied heavily on organophosphate insecticides (OPs) until the mid 2000s, when they were replaced with *Bacillus thuringiensis*. Resistance by southern French populations of *C. pipiens* to OPs involves both mutations that render the OP target, acetyl cholinesterase, less sensitive to the insecticide, and that increase expression of detoxifying

Figure 1



Schematic conceptualization of the effects of rate of migration from (a) a hotspot of resistance, or (b) a wild-type susceptible population, into receiving populations of high or low selection for resistance on relative rates of increase in resistance allele frequency (R freq) over time (generations). The dashed arc indicates the region around the resistance hotspot within which mitigation tactics must be implemented to slow the spread of resistance alleles through the larger landscape. Determining the spatial scale at which mitigation will effectively contain spread of resistance from a hotspot depends in part on the insect's dispersal rate and distance, and on spatial variation in selection pressure across the landscape. In practice, this scale can be very difficult to determine. If it extends beyond the dimensions of a farm, implementation of coordinated mitigation tactics by growers across an area may be necessary, with all the difficulties that implies. Thus, rapid implementation of mitigation around a still-localized hotspot offers the best hope of containment (see Andow *et al.* [22\*] for discussion in the case of *D. v. virgifera*).

esterases that degrade the insecticide. Both forms of resistance entail significant fitness costs in the absence of insecticide. The interplay of gene flow mediated by mosquito dispersal, selection for resistance near the coast, and fitness costs further inland produces clines in resistance allele frequencies. A joint analysis of the clines in target-site insensitivity and esterase overproduction produced an estimate of the standard deviation in parent-offspring dispersal of 6.6 km [3]. This rate of dispersal was sufficient to rapidly reestablish selection-migration equilibrium each year at both loci [4]. In the case of the *Ester*<sup>2</sup> locus in southern France, the *Ester*<sup>2</sup> resistance allele increased in frequency between 1999 and 2002 but did not replace *Ester*<sup>4</sup> despite a higher level of resistance, because it also imposed a higher fitness cost in the absence of insecticide [5]. After the use of OPs was discontinued in 2007, the *Ester*<sup>2</sup> allele was rapidly lost from the population. The *Ester*<sup>4</sup> allele persisted but the cline in *Ester*<sup>4</sup> frequency, while still significant, flattened markedly [6], presumably due to a combination of reduced selection at the coast and gene flow between the coast and inland.

#### Case study: spread of *Diabrotica virgifera virgifera* rotation resistance

Resistance to crop rotation in the western corn rootworm, *Diabrotica virgifera virgifera*, provided an opportunity to observe resistance spreading over a wide area as it occurred. Thus, the spatiotemporal dynamics of resistance were observed at a much larger geographical scale relative to adult dispersal distance than in the case of *C. pipiens* described above, so that alleles were not at migration-selection equilibrium during their years-long invasion of the surrounding metapopulation. The basic biology and history of adaptation to crop rotation by *D. v. virgifera* was comprehensively reviewed by Gray *et al.* [7]. Resistance to crop rotation is almost certainly based on a reduced preference for cornfields by females as oviposition sites. Thus in landscapes dominated by corn-soybean rotated crop fields, eggs (the overwintering stage) laid outside of corn have a better chance of hatching in a cornfield the following year than those that were laid in corn. In contrast to OP resistance in *C. pipiens*, the precise genetic basis of adaptation to crop rotation is not known. Recent research has focused on adaptations that allow

rotation resistant adult *D. v. virgifera* to cope with feeding on soybean foliage defended by plant protease inhibitors. Rotation resistant *D. v. virgifera* have elevated capthep-sin-L protease activity, compared to wild-type beetles [8]. This difference is related to differences in the gut flora of rotation resistant and susceptible beetles [9] which, in turn, may be related to differences in the expression of antimicrobial and immune-related genes [10,11]. It is currently unknown if these differences are part of the proximate mechanism of resistance to crop rotation or, more likely, a secondary adaptation improving fitness of beetles that encounter more soybean in their environment due to their decreased fidelity to cornfields.

Adaptation to crop rotation spread steadily from the original 1987 focus in Piper County, Illinois over the subsequent two decades. By 2007 populations throughout most of Illinois and neighboring Indiana were adapted to crop rotation, as were some populations in adjacent states. Little to no general genetic differentiation was observed between rotation resistant populations and surrounding susceptible populations [12,13]. This, in combination with the observed pattern of spread suggests that resistance to crop rotation moved across the landscape as a result of gene flow among established populations. Resistance to crop rotation apparently ceased to spread about a decade ago, after most of the region in which corn-soybean rotation is extremely common was occupied. Ovipositing in non-corn environments becomes more costly as the probability that such environments will be rotated to corn the following year declines. Furthermore, the advantage of the host-selective wild type behavior increases as the presence of corn becomes a more reliable indicator that corn will be present the following year (*i.e.*, continuous planting). Simulation studies suggest that the evolution and spread of this adaptation is impeded by increasing diversity of vegetation in the landscape [14,15]. A decrease in crop diversity in favor of continuously planted corn could be enough to impede spread of rotation resistance.

In recent years, the region in which rotation resistance is a problem appears to have receded (*e.g.*, [16]). This may be related to the commercialization of rootworm-targeting transgenic corn expressing various insecticidal proteins derived from *Bacillus thuringiensis* (Bt). Growers tend to plant rootworm Bt corn even in rotated fields, either to protect against rotation resistant populations, or because elite high-yielding non-Bt hybrids are difficult to obtain [17]. This practice reduces the selective advantage to rotation resistant phenotypes in rotated fields, because mortality is as high as among susceptible phenotypes ovipositing in non-rotated Bt corn [18]. Modeling indicates that the prevalence of rotation resistance in the landscape can even decrease if growers plant Bt corn more frequently in rotated fields than continuous fields [19].

#### Box 1 Larval dispersal, premating adult dispersal, and efficacy of Bt refuge configurations.

The most common type of proactive IRM strategy for Bt crops involves planting a non-Bt variety – the ‘refuge’ – as a nursery of susceptible insects. Refuges are often planted as blocks or strips of the required spatial dimensions in or near the larger Bt fields. A recent trend in some crop systems is to mix the requisite percentage of non-Bt refuge seed in a bag with the Bt seed. There is ongoing concern about the effectiveness of this blended refuge strategy in slowing resistance evolution. Larval dispersal between Bt and non-Bt plants can result in a sublethal dose, promoting differential survival of heterozygotes over homozygous susceptible larvae, thus accelerating accumulation of resistance alleles in the population. Models generally affirm that larval dispersal leads to faster evolution of resistance in a blended than in a block refuge system [17,50], unless substantial (but, unfortunately, realistic) grower noncompliance is included for the latter [51]. Recent research indicates that larval movement in several lepidopteran pest species is a threat to the blended refuge strategy [52–54]. Although less extensive than lepidopteran caterpillars, dispersal rate of subterranean rootworm larvae between adjacent plants is high enough [55] to potentially compromise a blended refuge strategy, as observed in laboratory evolution experiments [56]. However, in the case of *D. v. virgifera*, a blended refuge has the possible offsetting advantage of promoting mating of refuge adults with adults emerging from Bt plants. Most *D. v. virgifera* females mate near their natal plant, which can increase assortative mating among beetles emerging within Bt and refuge blocks, thus compromising the block refuge strategy [22,57]. Improved mixing of resistant and susceptible adults is often considered an advantage of blended refuge, which can be true for other species with limited premating dispersal [58], such as *L. decemlineata* [24]. But the argument of improved mixing of adults in a blended refuge field does not apply to species where significant dispersal from the natal plant occurs before mating. For example, young European corn borer, *Ostrinia nubilalis*, females disperse long distances before mating [59], a behavior common among many Lepidoptera, especially migratory species.

Understanding the evolutionary dynamics of rotation resistance in *D. v. virgifera* is important because crop rotation is the favored tactic for mitigating the emerging problem of resistance to Bt corn [20,21]. How fast and how far Bt resistance may spread is of concern to farmers who must manage the risk of crop losses caused by a product that could start to fail. Mitigation tactics are most effective while resistance is still spatially localized [22,23]. For monophagous or oligophagous insects with low premating dispersal, like *D. v. virgifera* and the Colorado potato beetle, *Leptinotarsa decemlineata*, crop rotation can be an excellent tactic to both limit spread of a resistance allele and lower its frequency *in situ* [22,24]. Rate of premating dispersal also impacts efficacy of various types of IRM refuges (Box 1).

#### Spread of resistance during species invasion or range expansion

When insects disperse into new territory, their initial population sizes are often small. Species introduced into an entirely new territory can experience severe bottlenecks [25,26], as can ‘pioneer’ populations of a species that is expanding its existing range [27]. Small

populations at the front of a range expansion can result in ‘gene surfing’, whereby rare alleles and/or new mutations are catapulted to high frequency by drift at the front and are then carried forward as the front moves, generating allele frequency clines [28,29]. Even mildly deleterious alleles can surf to high frequency [30], presumably including resistance alleles with moderate fitness costs.

*D. v. virgifera* and *L. decemlineata*, two chrysomelid pests notorious for evolving resistance, have undergone both range expansions and introductions to new continents. For both species, the likely ancestral range was in Mexico, but significant pest populations first arose in the Plains regions of Nebraska and Kansas, followed by eastward expansion across the United States and Canada. Both species were also subsequently introduced into Europe. Resistance to cyclodienes in *D. v. virgifera* is due to a mutation in the target gamma-amino butyric acid receptor gene, which exhibits a cline of increasing frequency toward the east coast of the United States [31]. This is surprising given that cyclodienes were withdrawn from use over a decade before the species reached the east coast and may represent an example of gene surfing [31]. Unlike populations in the United States, introduced European populations of *D. v. virgifera* are uniformly resistant to cyclodienes and susceptible to organophosphates [32]. Neither class of insecticide has been used extensively to control *D. v. virgifera* in Europe, so the current pattern of insecticide resistance there is probably a consequence of the North American populations that acted as the sources for introductions in Europe and bottlenecks during the introduction process [32]. The frequencies of an organophosphate resistance allele in *L. decemlineata* show a surprising spatial distribution. The allele is fixed in ancestral Mexican populations, is at high frequency (0.95) in US populations, and is variable, but lower on average, in Europe [33]. This suggests that the species may have been preadapted to organophosphates but lost resistance alleles during bottlenecks upon introduction to Europe.

### Increased resistance incidence across space: spread or independent evolution?

There is often uncertainty in the extent to which resistance evolves independently or spreads from a single source. Modeling indicates that the spatial scale over which resistance evolves depends in large part on the interplay between insect dispersal rate and the availability of refuge habitat in the landscape [60]. Multiple lines of evidence make it clear that *D. v. virgifera* resistance to various Bt toxins is evolving quickly and independently in local hotspots across a wide area [20,34,35]. Nevertheless, post-mating dispersal of *D. v. virgifera* can be extensive [36], and spatial spread from hotspots seems inevitable [22•]. Evidence suggests resistance spread from an unknown distance to fields in Iowa where damage to Cry34/35 Bt corn was detected [21]. The spatial

dimensions over which mitigation tactics must be employed to contain or eliminate a hotspot depends on a better understanding of long-range dispersal in this species, an area of active research [23•].

Codling moth, *Cydia pomonella*, dispersal seems to be mostly local, but with capacity for long-distance flights [37], causing uncertainty in whether regional variation in insecticide resistance is the result of independent evolution or spread [38]; most likely both processes are involved. Widespread resistance of *B. fusca* to Cry1Ab Bt corn developed rapidly after the crop’s introduction to South Africa, and was assumed to have evolved independently in different locations for widespread agronomic and refuge non-compliance reasons [39]. But, as explained by Peterson *et al.* [40], this is a pest with high dispersal capacity, and resistance could have spread quickly from a single location, especially given it is not a recessive trait. Population genetics analyses suggest that both phenomena are at work [41•]. Even in extreme cases, such as independent evolution of pyrethroid resistance in the Chagas’ disease vector, *Triatoma infestans*, at a micro-scale – for example, between neighboring dwellings – local spread from a focal point also seems to occur [42].

*S. frugiperda* is a complicated case where the relative roles of spread of Bt resistance by migration or independent evolution in hotspots remains uncertain. Cry1F resistance was detected in Puerto Rico in 2006 [43], and has since been found in Brazil [44] and parts of Florida [45]. *S. frugiperda* is a strong migrant, and the spread of Cry1F resistance from Puerto Rico to Florida by dispersing adults is possible. On the other hand, recent susceptibility surveys suggest resistance in different areas of Florida may be of independent evolutionary origin [46,47]. Resistance alleles were detected at low frequency in North Carolina, which receives annual migrants from Florida [48]. Population genetic and atmospheric trajectory analyses have effectively ruled out a Caribbean origin of the Cry1F resistance found in Brazil [49•].

Overall, theory and available data suggest that spatial and temporal distribution of resistance alleles differs depending on whether insect dispersal is spreading them among established populations or into new territory. In the former case, deterministic processes driven by selection, fitness costs and dominance appear to be central, whereas in the latter case, stochastic processes play a key role. One could argue that annual colonization of an agricultural field, or colonization by refuge insects of a Bt field cleared of conspecifics by the toxin, are cases of invasion on an accelerated temporal and micro-spatial scale. This seems analogous to the situation with *C. pipiens* mosquitoes in southern France, where resistance alleles annually come to migration-selection equilibrium at a small geographic scale relative to



individual dispersal distances. Comparing the dynamics of resistance evolution or spread in the case of classical invasion versus annual colonization of crops could be instructive for both. Large scale planting of Bt crops in concert with refuges to try and manage resistance evolution offers an exceptional opportunity to conduct such studies in a number of pest-crop systems.

## References and recommended reading

Papers of particular interest, published within the period of review, have been highlighted as:

- of special interest
- of outstanding interest

1. Sisterson MS, Carrière Y, Dennehy TJ, Tabashnik BE: **Evolution of resistance to transgenic crops: interactions between insect movement and field distribution.** *J. Econ. Entomol.* 2005, **98**:1751-1762.
  2. Alyokhin A, Miller L: **Propensity for flying and walking by the Colorado potato beetles treated with Imidacloprid.** *Am. J. Potato Res.* 2015, **92**:138-142.
  3. Lenormand T, Guillemaud T, Bourguet D, Raymond M: **Evaluating gene flow using selected markers: a case study.** *Genetics* 1998, **149**:1383-1392.
  4. Lenormand T, Bourguet D, Guillemaud T, Raymond M: **Tracking the evolution of insecticide resistance in the mosquito *Culex pipiens*.** *Nature* 1999, **400**:861-864.
  5. Labbé P, Sidos N, Raymond M, Lenormand T: **Resistance gene replacement in the mosquito *Culex pipiens*: fitness estimation from long-term cline series.** *Genetics* 2009, **182**:303-312.
  6. Milesi P, Lenormand T, Lagneau C, Weill M, Labbé P: **Relating fitness to long-term environmental variations in natura.** *Mol. Ecol.* 2016, **25**:5483-5499.
  7. Gray ME, Sappington TW, Miller NJ, Moeser J, Bohn MO: **Adaptation and invasiveness of western corn rootworm: intensifying research on an a worsening pest.** *Annu. Rev. Entomol.* 2009, **54**:303-321.
  8. Curzi MJ, Zavala JA, Spencer JL, Seufferheld MJ: **Abnormally high digestive enzyme activity and gene expression explain the contemporary evolution of a *Diabrotica* biotype able to feed on soybeans.** *Ecol. Evol.* 2012, **2**:2005-2017.
  9. Chu C-C, Spencer JL, Curzi MJ, Zavala JA, Seufferheld MJ: **Gut bacteria facilitate adaptation to crop rotation in the western corn rootworm.** *Proc. Natl. Acad. Sci.* 2013, **110**:11917-11922.
  10. Knolhoff LM, Walden KKO, Ratcliffe ST, Onstad DW, Robertson HM: **Microarray analysis yields candidate markers for rotation resistance in the western corn rootworm beetle, *Diabrotica virgifera virgifera*.** *Evol. Appl.* 2010, **3**:17-27.
  11. Chu C-C, Zavala JA, Spencer JL, Curzi MJ, Fields CJ, Drnevich J, Siegfried BD, Seufferheld MJ: **Patterns of differential gene expression in adult rotation-resistant and wild-type western corn rootworm digestive tracts.** *Evol. Appl.* 2015, **8**:692-704.
  12. Miller NJ, Kim KS, Ratcliffe ST, Estoup A, Bourguet D, Guillemaud T: **Absence of genetic divergence between western corn rootworms (Coleoptera: Chrysomelidae) resistant and susceptible to control by crop rotation.** *J. Econ. Entomol.* 2006, **99**:685-690.
  13. Miller NJ, Ciosi M, Sappington TW, Ratcliffe ST, Spencer JL, Guillemaud T: **Genome scan of *Diabrotica virgifera virgifera* for genetic variation associated with crop rotation tolerance.** *J. Appl. Entomol.* 2007, **131**:378-385.
  14. Onstad DW, Spencer JL, Guse CA, Levine E, Isard SA: **Modeling evolution of behavioral resistance by an insect to crop rotation.** *Entomol. Exp. Appl.* 2001, **100**:195-201.
  15. Onstad DW, Crowder DW, Isard SA, Levine E, Spencer JL, O'Neal ME, Ratcliffe ST, Gray ME, Bledsoe LW, DiFonzo C *et al.*: **Does landscape diversity slow the spread of rotation-resistant western corn rootworm (Coleoptera: Chrysomelidae)?** *Environ. Entomol.* 2003, **32**:992-1001.
  16. Dunbar MW, Gassmann AJ: **Abundance and distribution of western and northern corn rootworm (*Diabrotica* spp.) and prevalence of rotation resistance in eastern Iowa.** *J. Econ. Entomol.* 2013, **106**:168-180.
  17. Onstad DW, Mitchell PD, Hurley TM, Lundgren JG, Porter RP, Krupke CH, Spencer JL, DiFonzo CD, Baute TS, Hellmich RL *et al.*: **Seeds of change: corn seed mixtures for resistance management and integrated pest management.** *J. Econ. Entomol.* 2011, **104**:343-352.
  18. Tinsley NA, Spencer JL, Estes RE, Prasifka JR, Schrader PM, French BW, Gray ME: **Larval mortality and development for rotation-resistant and rotation-susceptible populations of western corn rootworm on Bt corn.** *J. Appl. Entomol.* 2015, **139**:46-54.
- Demonstrates that mortality of rotation resistant and wild type susceptible *D. v. virgifera* larvae are equally susceptible to Cry3Bb1 and Cry34/35Ab1 Bt toxins, singly and when pyramided. This key finding supports the idea that the halt of geographic spread, and indeed apparent reversal, of rotation resistance was caused partly or wholly by high adoption of Bt corn for rotated fields.
19. Crowder DW, Onstad DW, Gray ME, Pierce CMF, Hager AG, Ratcliffe ST, Steffey KL: **Analysis of the dynamics of adaptation to transgenic corn and crop rotation by western corn rootworm (Coleoptera: Chrysomelidae) using a daily time-step model.** *J. Econ. Entomol.* 2005, **98**:534-551.
  20. Gassmann AJ, Petzold-Maxwell JL, Clifton EH, Dunbar MW, Hoffmann AM, Ingber DA, Keweshan RS: **Field-evolved resistance by western corn rootworm to multiple *Bacillus thuringiensis* toxins in transgenic maize.** *Proc. Natl. Acad. Sci.* 2014 <http://dx.doi.org/10.1073/pnas.1317179111>.
  21. Gassmann AJ, Shrestha RB, Jakka SRK, Dunbar MW, Clifton EH, Paolino AR, Ingber DA, French BW, Masloski KE, Dounda JW *et al.*: **Evidence of resistance to Cry34/35 Ab1 corn by western corn rootworm (Coleoptera: Chrysomelidae): root injury in the field and larval survival in plant-based bioassays.** *J. Econ. Entomol.* 2016, **109**:1872-1880.
  22. Andow DA, Pueppke SG, Schaafsma AW, Gassmann AJ, Sappington TW, Meinke LJ, Mitchell PD, Hurley TM, Hellmich RL, Porter RP: **Early detection and mitigation of resistance to Bt maize by western corn rootworm (Coleoptera: Chrysomelidae).** *J. Econ. Entomol.* 2016, **109**:1-12.
- Analyzes the various causes of rapid evolution of resistance in *D. v. virgifera* to Bt corn in the field, including evidence for both *de novo* evolution and spread from hotspots. Recommends crop rotation as the most effective tactic for mitigating resistance. Discusses the need to better understand rootworm dispersal to determine the geographic scale for implementing mitigation tactics to spatially contain resistance in areas where it is detected.
23. Martinez JC, Caprio MA: **IPM use with the deployment of a non-high dose Bt pyramid and mitigation of resistance for western corn rootworm (*Diabrotica virgifera virgifera*).** *Environ. Entomol.* 2016, **45**:747-761.
- Model simulations of various tactics to mitigate or spatially contain *D. v. virgifera* resistance to Bt corn showed that crop rotation was a very good option, but must be implemented quickly after resistance develops to be effective. Also showed that mitigation of a resistance hotspot must extend spatially beyond the hotspot itself to contain spread by dispersing insects.
24. Huseth AS, Petersen JD, Poveda K, Szendrei Z, Nault BA, Kennedy GG, Groves RL: **Spatial and temporal potato intensification drives insecticide resistance in the specialist herbivore, *Leptinotarsa decemlineata*.** *PLoS One* 2015, **10**: e0127576.
  25. Ciosi M, Miller NJ, Kim KS, Giordano R, Estoup A, Guillemaud T: **Invasion of Europe by the western corn rootworm, *Diabrotica virgifera virgifera*: multiple transatlantic introductions with various reductions of genetic diversity.** *Mol. Ecol.* 2008, **17**:3614-3627.

26. Miller NJ, Estoup A, Toepfer S, Bourguet D, Lapchin L, Derridj S, Kim KS, Reynaud P, Furlan L, Guillemaud T: **Multiple transatlantic introductions of the western corn rootworm.** *Science* 2005, **310**:992.
27. Excoffier L, Ray N: **Surfing during population expansions promotes genetic revolutions and structuration.** *Trends Ecol. Evol.* 2008, **23**:347-351.
28. Hallatschek O, Nelson DR: **Gene surfing in expanding populations.** *Theor. Popul. Biol.* 2008, **73**:158-170.
29. Edmonds CA, Lillie AS, Cavalli-Sforza LL: **Mutations arising in the wave front of an expanding population.** *Proc. Natl. Acad. Sci. U. S. A.* 2004, **101**:975-979.
30. Travis JMJ, Münkemüller T, Burton OJ, Best A, Dytham C, Johst K: **Deleterious mutations can surf to high densities on the wave front of an expanding population.** *Mol. Biol. Evol.* 2007, **24**:2334-2343.
31. Wang H, Coates BS, Chen H, Sappington TW, Guillemaud T, Siegfried BD: **Role of a gamma-aminobutyric acid (GABA) receptor mutation in the evolution and spread of *Diabrotica virgifera virgifera* resistance to cyclodiene insecticides.** *Insect Mol. Biol.* 2013, **22**:473-484.
32. Ciosi M, Toepfer S, Li H, Haye T, Kuhlmann U, Wang H, Siegfried B, Guillemaud T: **European populations of *Diabrotica virgifera virgifera* are resistant to aldrin, but not to methyl-parathion.** *J. Appl. Entomol.* 2009, **133**:307-314.
33. Piironen S, Lindström L, Lyytinen A, Mappes J, Chen YH, Izzo V, Grapputo A: **Pre-invasion history and demography shape the genetic variation in the insecticide resistance-related acetylcholinesterase 2 gene in the invasive Colorado potato beetle.** *BMC Evol. Biol.* 2013, **13**:13.
34. Ingber DA, Gassmann AJ: **Inheritance and fitness costs of resistance to Cry3 Bb1 corn by western corn rootworm (Coleoptera: Chrysomelidae).** *J. Econ. Entomol.* 2015, **108**:2421-2432.
35. Wangila DS, Gassmann AJ, Petzold-Maxwell JL, French BW, Meinke LJ: **Susceptibility of Nebraska western corn rootworm (Coleoptera: Chrysomelidae) populations to Bt corn events.** *J. Econ. Entomol.* 2015, **108**:742-751.
36. Meinke LJ, Sappington TW, Onstad DW, Guillemaud T, Miller NJ, Komaromi J, Levay N, Furlan L, Kiss J, Toth F: **Western corn rootworm (*Diabrotica virgifera virgifera* LeConte) population dynamics.** *Agric. For. Entomol.* 2009, **11**:29-46.
37. Margaritopoulos JT, Voudouris CC, Olivares J, Sauphanor B, Mamuris Z, Tsitsipis JA, Franck P: **Dispersal ability in codling moth: mark-release-recapture experiments and kinship analysis.** *Agric. For. Entomol.* 2012, **14**:399-407.
38. Grigg-McGuffin K, Scott IM, Bellerose S, Chouinard G, Cormier D, Scott-Dupree C: **Susceptibility in field populations of codling moth, *Cydia pomonella* (L.) (Lepidoptera: Tortricidae), in Ontario and Quebec apple orchards to a selection of insecticides.** *Pest Manag. Sci.* 2015, **71**:234-242.
39. Kruger M, Van Rensburg JBJ, Van den Berg J: **Perspective on the development of stem borer resistance to Bt maize and refuge compliance at the Vaalharts irrigation scheme in South Africa.** *Crop Prot.* 2009, **28**:684-689.
40. Peterson B, Bezuidenhout CC, Van den Berg J: **Cytochrome c oxidase I and cytochrome b gene sequences indicate low genetic diversity in South African *Busseola fusca* (Lepidoptera: Noctuidae) from maize.** *Afr. Entomol.* 2016, **24**:518-523.
41. Campagne P, Capdevielle-Dulac C, Pasquet R, Cornell SJ, Kruger M, Silvain J-F, LeRü B, Van den Berg J: **Genetic hitchhiking and resistance evolution to transgenic Bt toxins: insights from the African stalk borer *Busseola fusca* (Noctuidae).** *Heredity* 2017, **118**:330-339.
42. Mougabure-Cueto G, Picollo MI: **Insecticide resistance in vector Chagas disease: evolution, mechanisms and management.** *Acta Trop.* 2015, **149**:70-85.
43. Storer NP, Babcock JM, Schlenz M, Meade T, Thompson GD, Bing JW, Huckaba RM: **Discovery and characterization of field resistance to Bt maize: *Spodoptera frugiperda* (Lepidoptera: Noctuidae) in Puerto Rico.** *J. Econ. Entomol.* 2010, **103**:1031-1038.
44. Farias JR, Andow DA, Horikoshi RJ, Sorgatto RJ, Fresia P, dos Santos AC, Omoto C: **Field-evolved resistance to Cry1F maize by *Spodoptera frugiperda* (Lepidoptera: Noctuidae) in Brazil.** *Crop Prot.* 2014, **64**:150-158.
45. Huang F, Qureshi JA, Meagher RL, Reising DD, Head GP, Andow DA, Ni X, Kerns D, Buntin GD, Niu Y *et al.*: **Cry1F resistance in fall armyworm *Spodoptera frugiperda*: single gene versus pyramided Bt maize.** *PLoS One* 2014, **9**:e112958.
46. Dangal V, Huang F: **Fitness costs of Cry1F resistance in two populations of fall armyworm, *Spodoptera frugiperda* (JE Smith), collected from Puerto Rico and Florida.** *J. Invertebr. Pathol.* 2015, **127**:81-86.
47. Da Silva KF, Spencer TA, Crespo ALB, Siegfried BD: **Susceptibility of *Spodoptera frugiperda* (Lepidoptera: Noctuidae) field populations to the Cry1F *Bacillus thuringiensis* insecticidal protein.** *Fla. Entomol.* 2016, **99**:629-633.
48. Li G, Reising D, Miao J, Gould F, Huang F, Feng H: **Frequency of Cry1F non-recessive resistance alleles in North Carolina field populations of *Spodoptera frugiperda* (Lepidoptera: Noctuidae).** *PLoS One* 2016, **11**:e0154492.
49. Nagoshi RN, Fleischer S, Meagher RL, Hay-Roe M, Khan A, Murua MG, Silvie P, Vergara C, Westbrook J: **Fall armyworm migration across the Lesser Antilles and the potential for genetic exchanges between North and South American populations.** *PLoS One* 2017, **12**:e0171743.
- Combined genetic markers and HYSPLIT air transport trajectories to demonstrate that there is little likelihood of migration of Cry1 F-resistant *S. frugiperda* from Puerto Rico to southern Caribbean islands or South America. This group has conducted very interesting and rigorous research on migratory routes of this species in the Americas with far-reaching implications for pest management and resistance spread, and the Introduction provides a good synopsis of this work.
50. Garcia AG, Ferreira CP, Consoli FL, Godoy WAC: **Predicting evolution of insect resistance to transgenic crops in within-field refuge configurations, based on larval movement.** *Ecol. Complex.* 2016, **28**:94-103.
51. Pan Z, Onstad D, Crain P, Crespo A, Hutchison W, Buntin D, Porter P, Catchot A, Cook D, Pilcher C *et al.*: **Evolution of resistance by *Helicoverpa zea* (Lepidoptera: Noctuidae) infesting insecticidal crops in the Southern United States.** *J. Econ. Entomol.* 2016, **109**:821-831.
- Model simulations provide an appreciation for the great complexity of resistance evolution in a pest like *H. zea* that is under selection pressure from both Bt cotton and Bt corn in the agricultural landscape of the southern US. They recognize that long distance migration, which they did not try to include in their models, could add an even greater level of complexity to the situation. Results also showed that blended refuges perform better than block refuges in delaying resistance only in the context of low grower refuge compliance.
52. Erasmus A, Marais J, Van den Berg J: **Movement and survival of *Busseola fusca* (Lepidoptera: Noctuidae) larvae within maize plantings with different ratios of non-Bt and Bt seed.** *Pest Manag. Sci.* 2016, **72**:2287-2294.
53. Miraldo LL, Bernardi O, Horikoshi RJ, Amaral FSA, Bernardi D, Omoto C: **Functional dominance of different aged larvae of Bt-resistant *Spodoptera frugiperda* (Lepidoptera: Noctuidae) on transgenic maize expressing Vip3 Aa20 protein.** *Crop Prot.* 2016, **88**:65-71.
54. Pannuti LER, Paula-Moraes SV, Hunt TE, Baldin ELL, Dana L, Malaquias JV: **Plant-to-plant movement of *Striacosta albicosta* (Lepidoptera: Noctuidae) and *Spodoptera frugiperda* (Lepidoptera: Noctuidae) in Maize (*Zea mays*).** *J. Econ. Entomol.* 2016, **109**:1125-1131.

55. Zukoff SN, Zukoff AL, Geisert RW, Hibbard BE: **Western corn rootworm (Coleoptera: Chrysomelidae) larval movement in eCry3.1Ab+mCry3A seed blend scenarios.** *J. Econ. Entomol.* 2016, **109**:1834-1845.
56. Deitloff J, Dunbar MW, Ingber DA, Hibbard BE, Gassmann AJ: **Effects of refuges on the evolution of resistance to transgenic corn by the western corn rootworm, *Diabrotica virgifera virgifera* LeConte.** *Pest Manag. Sci.* 2016, **72**:190-198.
57. Hughson SA, Spencer JL: **Emergence and abundance of western corn rootworm (Coleoptera: Chrysomelidae) in Bt cornfields with structured and seed blend refuges.** *J. Econ. Entomol.* 2015, **108**:114-125.
- Field experiments examining in exquisite detail and fine-grained spatial scale adult *D. v. virgifera* dispersal to and from different Bt corn refuge configurations throughout the season. The results show that the assumption of random mating of rootworm adults emerging from a Bt field with those from refuge is most likely to be met with a blended refuge strategy rather than block refuge.
58. Campagne P, Smouse PE, Pasquet R, Silvain J-F, Le Ru B, Van den Berg J: **Impact of violated high-dose refuge assumptions on evolution of Bt resistance.** *Evol. Appl.* 2016, **9**:596-607.
- Uses modeling to explore the consequences of various violations of the high-dose refuge strategy of IRM for resistance evolution. Of special relevance to our review are results and discussion of various adult dispersal scenarios under different kinds of refuge configurations: several can lead to increased assortative mating of resistant and susceptible individuals, which is a good way to accelerate resistance evolution.
59. Dorhout DL, Sappington TW, Rice ME: **Evidence for obligate migratory flight behavior in young European corn borer (Lepidoptera: Crambidae) females.** *Environ. Entomol.* 2008, **37**:1280-1290.
60. Huang Y, Qin Y, Feng H, Wan P, Li Z: **Modeling the evolution of insect resistance to one- and two-toxin Bt-crops in spatially heterogeneous environments.** *Ecol. Modell.* 2017, **347**:72-84.