



Complete Genome Sequence of *Campylobacter hepaticus* USA52, Associated with Chicken Spotty Liver Disease

 Zuowei Wu,^a Orhan Sahin,^b Qijing Zhang^a

^aVeterinary Microbiology and Preventive Medicine, Iowa State University, Ames, Iowa, USA

^bVeterinary Diagnostic and Production Animal Medicine, Iowa State University, Ames, Iowa, USA

ABSTRACT *Campylobacter hepaticus* was recently identified as the etiological agent of chicken spotty liver disease. Here, we report the complete genome sequence of *C. hepaticus* strain USA52 from the United States. The genome comprises a chromosome of 1,509,100 bp with an average GC content of 28.02%.

Spotty liver disease (SLD) is an acute infectious disease of layer poultry and has become an important concern for the poultry egg and meat industries (1–3). Although the disease was originally reported in the United States and Canada in the 1950s, the definitive etiological agent was not identified until 2015, when a new *Campylobacter* species, *C. hepaticus*, was determined to be the causative agent in the United Kingdom and Australia (4–6). In the United States, *C. hepaticus* was first identified from SLD layers in 2017 (7). However, to date, a complete genome sequence of *C. hepaticus* from the United States has not been reported. Here, we report the complete genome sequence of *C. hepaticus* strain USA52.

C. hepaticus USA52 was isolated from the bile of a dead SLD layer as described previously (4). The bile sample was submitted to our diagnostic lab by a farm with an SLD outbreak in Nebraska. Strain USA52 was identified to be *C. hepaticus* by 16S rRNA gene sequencing and PCR detection of the glycerol kinase gene as described previously (7). Strain USA52 cells were collected after culturing on blood agar plates (Remel, Lenexa, KS) for 3 days at 37°C, and genomic DNA was extracted using a Genomic-tip 20/G kit (Qiagen, Germantown, MD). A Nanopore library was constructed with 1.5 μg of genomic DNA using the Nanopore ligation sequencing kit (SQK-LSK108) (Oxford Nanopore Technologies, Cambridge, MA). Short DNA fragments (<10 kb) were removed from the library using a short-read eliminator (SRE) XS kit (Circulomics, Baltimore, MD). The library was sequenced on the Oxford Nanopore GridION X5 sequencing platform (flow cell, FLO-MIN106), and the reads were base called using Guppy v3.2.8 (8) at Iowa State University. The library yielded 4,000 reads with an N_{50} value of 19,540 bp, providing 394-fold coverage for the complete genome sequence. An Illumina short-read library was prepared with 1 μg of the same genomic DNA extract as above using the NEBNext Ultra II DNA FS library kit (New England Biolabs, Ipswich, MA) and sequenced on the Illumina MiSeq platform in 150-bp paired-end mode at Iowa State University, yielding 99-fold genome coverage with 1,003,036 reads. The read quality was examined using FastQC v0.11.9 (9). The Nanopore reads were *de novo* assembled using Flye v2.4 (10), producing one 1,559,990-bp contig. The contig was then error corrected with the Illumina reads using Pilon v1.23 (11). The corrected contig was circularized into a chromosome of 1,509,100 bp with a GC content of 28.02% by removing the overlapping ends manually after aligning it using Mauve v2.13 (12), with the contigs (assembled by Velvet v1.2.10 [13]) from the Illumina reads. The genes were annotated using the NCBI Prokaryotic Genome Annotation Pipeline (PGAP) v4.13 (14). The chromosome sequence was rotated to start at the *dnaA* gene using GAMOLA2 (15). Default parameters were applied for all software. The chromosome contained 1,484 predicted

Citation Wu Z, Sahin O, Zhang Q. 2021. Complete genome sequence of *Campylobacter hepaticus* USA52, associated with chicken spotty liver disease. Microbiol Resour Announc 10:e01266-20. <https://doi.org/10.1128/MRA.01266-20>.

Editor Vincent Bruno, University of Maryland School of Medicine

Copyright © 2021 Wu et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to Zuowei Wu, wuzw@iastate.edu.

Received 4 November 2020

Accepted 6 February 2021

Published 25 February 2021

protein-coding sequences (CDSs), 3 sets of rRNA genes, 3 noncoding RNA (ncRNA) genes, and 43 tRNA genes.

No reviews from the Institutional Ethics Committee and the Institutional Animal Care and Use Committee were required, because this study did not include any animal studies.

Data availability. The complete genome sequence of *C. hepaticus* USA52 has been deposited at GenBank under accession number [CP063536](https://doi.org/10.1186/s13059-019-1727-y). The raw sequence data have been deposited at the GenBank Sequence Read Archive under accession numbers [SRR12900734](https://doi.org/10.1073/pnas.1604560113) (Illumina reads) and [SRR12900733](https://doi.org/10.1073/pnas.1604560113) (Nanopore reads).

ACKNOWLEDGMENT

This work was funded by the Iowa Livestock Health Advisory Council.

REFERENCES

- Crawshaw T, Young S. 2003. Increased mortality on a free-range layer site. *Vet Rec* 153:664.
- Burch D. 2005. Avian vibriotic hepatitis in laying hens. *Vet Rec* 157:528. <https://doi.org/10.1136/vr.157.17.528-a>.
- Grimes T, Reece R. 2011. Spotty liver disease—an emerging disease in free-range egg layers in Australia, p 53–56. *In* Proceedings of the 60th Western Poultry Disease Conference. American Association of Avian Pathologists, Jacksonville, FL.
- Crawshaw TR, Chanter JI, Young SCL, Cawthraw S, Whatmore AM, Koyle MS, Vidal AB, Salguero FJ, Irvine RM. 2015. Isolation of a novel thermophilic *Campylobacter* from cases of spotty liver disease in laying hens and experimental reproduction of infection and microscopic pathology. *Vet Microbiol* 179:315–321. <https://doi.org/10.1016/j.vetmic.2015.06.008>.
- Van TTH, Elshagmani E, Gor MC, Scott PC, Moore RJ. 2016. *Campylobacter hepaticus* sp. nov., isolated from chickens with spotty liver disease. *Int J Syst Evol Microbiol* 66:4518–4524. <https://doi.org/10.1099/ijsem.0.001383>.
- Van TTH, Elshagmani E, Gor M-C, Anwar A, Scott PC, Moore RJ. 2017. Induction of spotty liver disease in layer hens by infection with *Campylobacter hepaticus*. *Vet Microbiol* 199:85–90. <https://doi.org/10.1016/j.vetmic.2016.12.033>.
- Gregory M, Klein B, Sahin O, Girgis G. 2018. Isolation and characterization of *Campylobacter hepaticus* from layer chickens with spotty liver disease in the United States. *Avian Dis* 62:79–85. <https://doi.org/10.1637/11752-092017-Reg.1>.
- Wick RR, Judd LM, Holt KE. 2019. Performance of neural network basecalling tools for Oxford Nanopore sequencing. *Genome Biol* 20:129. <https://doi.org/10.1186/s13059-019-1727-y>.
- Andrews S. 2010. FastQC: a quality control tool for high throughput sequence data. <http://www.bioinformatics.babraham.ac.uk/projects/fastqc/>.
- Lin Y, Yuan J, Kolmogorov M, Shen MW, Chaisson M, Pevzner PA. 2016. Assembly of long error-prone reads using de Bruijn graphs. *Proc Natl Acad Sci U S A* 113:E8396–E8405. <https://doi.org/10.1073/pnas.1604560113>.
- Walker BJ, Abeel T, Shea T, Priest M, Abouelliel A, Sakthikumar S, Cuomo CA, Zeng Q, Wortman J, Young SK, Earl AM. 2014. Pilon: an integrated tool for comprehensive microbial variant detection and genome assembly improvement. *PLoS One* 9:e112963. <https://doi.org/10.1371/journal.pone.0112963>.
- Darling ACE, Mau B, Blattner FR, Perna NT. 2004. Mauve: multiple alignment of conserved genomic sequence with rearrangements. *Genome Res* 14:1394–1403. <https://doi.org/10.1101/gr.2289704>.
- Zerbino DR. 2010. Using the Velvet de novo assembler for short-read sequencing technologies. *Curr Protoc Bioinformatics* Chapter 11:Unit 11.15. <https://doi.org/10.1002/0471250953.bi1105s31>.
- Tatusova T, DiCuccio M, Badretdin A, Chetvermin V, Nawrocki EP, Zaslavsky L, Lomsadze A, Pruitt KD, Borodovsky M, Ostell J. 2016. NCBI Prokaryotic Genome Annotation Pipeline. *Nucleic Acids Res* 44:6614–6624. <https://doi.org/10.1093/nar/gkw569>.
- Altermann E, Lu J, McCulloch A. 2017. GAMOLA2, a comprehensive software package for the annotation and curation of draft and complete microbial genomes. *Front Microbiol* 8:346. <https://doi.org/10.3389/fmicb.2017.00346>.